arXiv:1203.2338v1 [math.AG] 11 Mar 2012

Irregular Hodge Filtration on Twisted De Rham Cohomology*

Jeng-Daw Yu†

Abstract

We define the irregular Hodge filtration on the exponentially twisted de Rham cohomology of a smooth quasi-projective complex variety. Basic properties are studied and we compare the filtration with a few others appeared in literature.

Introduction

(a) *The goal*

Let U be a complex smooth quasi-projective variety and $f \in \mathcal{O}(U)$ be a regular function on U. Consider the algebraic connection $\nabla = \nabla_f = d + df$ on the structure sheaf $\mathcal{O}$ of U defined by

$$\begin{array}{rcl} \nabla : \mathcal{O} & \to & \Omega^1 \\ v & \mapsto & dv + v \cdot df. \end{array}$$

It is clear that ∇ is integrable and hence extends to a chain map, still denoted by ∇, on the sheaves $\Omega^\bullet$ of differential forms of U. The hypercohomology of the complex $(\Omega^\bullet, \nabla)$ on U is by definition the de Rham cohomology $H_{\mathrm{dR}}(U, \nabla)$ of the connection ∇, which is a finite collection of finite dimensional complex vector spaces. When f is a constant, we recover the algebraic de Rham cohomology $H_{\mathrm{dR}}(U/\mathbb{C})$ of U, which is equipped with a Hodge filtration coming from various truncations $(\Omega^{\geq p}, d)$ of the usual de Rham complex.

(∇ is the *exponential twist* of the usual differential d in the sense that the diagram

$$\begin{tikzcd} \mathcal{O} \arrow[r, "\nabla"] \arrow[d, "\cong"'] & \Omega^1 \arrow[d, "\cong"] \\ \exp f \cdot \mathcal{O} \arrow[r, "d"] & \exp f \cdot \Omega^1 \end{tikzcd}$$

commutes. Here the vertical arrows are the multiplication by the exponential $\exp(f)$ of f. However since the function $\exp(f)$ is transcendental if f is non-trivial, one should regard the $\exp(f)$ in the lower corners as a symbol of a basis.)

---

[*] This work was partially supported by the Golden-Jade fellowship of Kenda Foundation, the NCTS (Main Office) and the NSC, Taiwan.

[†] Department of Mathematics, National Taiwan University, Taipei, Taiwan; `jdyu@math.ntu.edu.tw`.

When U is a curve, Deligne [9, pp.109-128], motivated by the analogues between algebraic connections with irregular singularities and lisse étale sheaves with wild ramifications, has defined an *irregular Hodge filtration* F^λ, indexed by $\lambda \in \mathbb{R}$, on $H_{\mathrm{dR}}(U, \nabla)$. More precisely, let X be the smooth compactifiaction of U and $S = X \setminus U$ the complement. The function f on U then extends to a rational function on X and hence ∇ defines a meromorphic connection

$$\nabla : \mathcal{O}_X(*S) \to \Omega_X^1(*S)$$

between functions and forms with poles supported on S. We have

$$H^i_{\mathrm{dR}}(U, \nabla) = \mathbb{H}^i\left(X, \mathcal{O}_X(*S) \xrightarrow{\nabla} \Omega^1_X(*S)\right). \tag{1}$$

Deligne then defines an exhaustive and separated decreasing filtration $F^\lambda(\nabla)$ on the above two-term complex. The desired irregular Hodge filtration on H_{dR} is then given by

$$F^\lambda H^i_{\mathrm{dR}}(U, \nabla) := \mathrm{Image}\left\{\mathbb{H}^i\left(X, F^\lambda(\nabla)\right) \to H^i_{\mathrm{dR}}(U, \nabla)\right\} \tag{2}$$

under the identification (1). When f is a constant, $F^\bullet(\nabla)$ reduces to the pole-order filtration $P^\bullet$ defined in [7, II.3.12] and thus one recovers the usual Hodge filtration.

Deligne has showed that the spectral sequence associated with this filtration degenerates at the initial stage (i.e. the arrow in (2) is always injective) and proved that the irregular Hodge filtration respects the pairing between the cohomology of ∇ and of the dual connection. However the filtration and its complex conjugate are not opposite to each other in general. We remark that in [9], the definition of the irregular Hodge filtration is justified by its relation with the expected weights of special values of the gamma function. Moreover the filtration is defined for more general connections of certain type and not necessarily of rank one.

In this paper, we propose a definition of the irregular Hodge filtration for ∇ on U of arbitrary dimension. The idea is similar to Deligne's approach. We first pick a compactification X of U such that f extends to a morphism from X to $\mathbb{P}^1$ and that the complement $S := X \setminus U$ is a normal crossing divisor. We then define a decreasing filtration $F^\lambda(\nabla)$ of the twisted meromorphic de Rham complex $(\Omega^\bullet_X(*S), \nabla)$. The sheaves involved in each $F^\lambda(\nabla)$ are all locally free on X. However the new definition does not coincide with Deligne's when U is a curve. In fact when f is constant, our filtration is not the same as the pole-order filtration $P^\bullet$ but is equal to the usual Hodge filtration $(\Omega_X^{\geq\lambda}(\log S), d)$ of the de Rham complex of logarithmic differential forms. Although the filtration fails to be exhaustive in general, it is rich enough to capture the de Rham cohomology of ∇ and indeed induces the same filtration on the cohomology as Deligne's in the curve case.

Another advantage of using the logarithmic differential forms lies in the fact that unlike the curve case, one does not have a canonical choice of the compact X. Two different choices are connected by a birational morphism π and the sheaves $\Omega^p_X(\log S)$ behave well under π. We shall prove that the irregular Hodge filtration on the de Rham cohomology of ∇ obtained in this way is independent of the choice of X and satisfies some functorial properties. As in the curve case, we also demonstrate that the filtration respects the Poincaré pairing between the cohomology of ∇ and of the dual connection.

Unfortunately at this point, we do not know when the associated Hodge to de Rham spectral sequence degenerates in general. However we provide several examples of relating this irregular Hodge filtration to other filtrations appeared in literature that known to admit the degeneracy of the spectral sequences. Some new techniques might be needed to attack the degeneracy problem. An ideal scene would be to obtain both analytic method and algebraic method of the proof if the degeneracy turns out to be true. For the recent developments of harmonic methods related to the irregular connections in more general settings, see the survey [26] of a 2012 Bourbaki talk. On the algebraic side, one may need to develop a mod p version of p-adic (e.g. rigid) cohomology theory which allows to have an irregular version of the Cartier isomorphism in order to adapt the method of [8]. It seems out of reach at this point.

To close the discussion here of the story in characteristic 0, we mention that in the direction of relating the algebraic de Rham cohomology to a Betti type cohomology attached to any integrable algebraic connection of arbitrary rank via periods, the homology with coefficients in rapid decay simplicial chains has been defined and the duality to the de Rham cohomology has been established in [6] for the curve case and [15] in general. On the other hand, the relation to the nonabelian Hodge theory has been discussed in [23]. The exponentially twisted de Rham cohomology also appears in the theory of mirror symmetry [18] and the study of Donaldson-Thomas invariants [19]. We hope the investigation of the irregular Hodge filtration can provide more structures and shed some light into these areas. In [25], another generalization of the irregular Hodge filtration in the higher rank case over a projective line is developed and has been connected to the so-called supersymmatric index. We will come back to the last mentioned paper in §6.

(b) *Motivation from exponential sums*

The other motivation of this work is to see if one can generalize Katz's conjecture to the p-adic estimates of exponential sums. Here the original conjecture refers to the relation between the Hodge numbers of a smooth projective X in positive characteristic and the p-adic divisibility of the absolute Frobenius acting on the crystalline cohomology of X. For the precise statements, see [5, Thm.8.26], [17, §II.4]; see [21] for generalizations to powers of Frobenius. The corresponding statements for more general coefficients with possibly non-trivial logarithmic structures can be found in [22, §7.3].

Now for a smooth quasi-projective U over a finite field κ and a regular function $f : U \to \mathbb{A}^1$, one defines its L-function $L(T)$ constructed from the exponential sums of f over various finite field extensions of κ (after choosing an additive character of κ). $L(T)$ is known to be a rational function whose zeros and poles are Weil numbers. (In fact the Lefschetz fixed point formula in either ℓ-adic étale cohomology or rigid cohomology tells us that $L(T)$ is a factor of the usual zeta-function of the Artin-Schreier cover of U associated with f.) In this setting, one hopes that the irregular Hodge numbers would provide a similar p-adic divisibility property of the Frobenius action and hence p-adic estimates of the L-function and the exponential sums.

This application of the irregular Hodge filtration still seems far from realizable in general even in the case where U is a curve. However when U is an open set of $\mathbb{P}^1$, it is known that the irregular Hodge numbers indeed provide a lower bound of the Newton polygon of $L(T)$ by the work [29]. (One checks that the Hodge numbers defined in that paper coincide with the irregular Hodge numbers for the corresponding connection over the complex numbers. Also see [9, second paragraph in p.175] where Deligne has noticed the validity in the case of Gauss sums.) Another known example is the case of exponential sums over a torus by the work of, e.g., [1, 27]. In this toric case, the irregular Hodge filtration is closely related to the Newton polyhedron filtration studied in [20, 2] and recalled in §5. If one ignores the Hodge filtrations, a version of the comparison between the rigid cohomology associated with exponential sums and the de Rham cohomology of the corresponding connections has been established in [3, Cor.3.8].

(c) *Back to the paper*

The main body of the paper is organized as follows. In §1 we give the definition of the irregular Hodge filtration of the twisted de Rham complex on a certain compactification X of U. We show that the induced filtration on the de Rham cohomology of (U, ∇) is independent of X. Along the way some basic properties of the filtration are derived. We shall define the corresponding filtration on the cohomology with compact support. We then establish the perfect Poincaré pairing between the de Rham cohomology of ∇ and of its dual with compact support in §2. The irregular Hodge filtrations on them are shown to respect the Poincaré pairing. In fact since we do not know when the Hodge to de Rham spectral sequence degenerates, we will also define pairings between terms on each stage of the spectral sequence and discuss their relations.

§3 to §6 are devoted to providing examples. In §3 we give a simple illustration of computing the irregular Hodge filtration of a certain connection on a regular surface. In §4 we discuss the case where $U = \mathbb{A}^1 \times U'$ and f is the direct product of the identity on $\mathbb{A}^1$ and a function f' on U'. In this case, the irregular Hodge filtration reduces to the usual Hodge filtration of the subvariety defined by f' if it is smooth. We then recall the work of Adolphson and Sperber on the twisted de Rham cohomology over a torus in §5. In this case a filtration coming from the Newton polyhedron Δ of the function f is defined and the associated spectral sequence is shown to degenerate if f is *non-degenerate* with respect to Δ in [2]. We show that in this case the filtration from Δ on the de Rham cohomology coincides with our irregular Hodge filtration. In §6 we recall the work [25] of Sabbah on the generalization of Deligne's irregular Hodge filtration and apply it to the study of the case where $f : X \to \mathbb{P}^1$ is smooth and $U = f^{-1}(\mathbb{A}^1)$.

Finally in the appendix we briefly recall Deligne's definition of the irregular Hodge filtration in the curve case and indicate that his definition gives the same filtration on the de Rham cohomology as ours.

Of course, the reader is referred to the main body for definitions, various assumptions and precise statements.

Part of this work has been done during my visit of Universität Duisburg - Essen in 2011 - 2012. I am grateful for the financial support, the hospitality and the inspiring environment provided by the group of *Essener Seminar für Algebraische Geometrie und Arithmetik*. I thank Professor Hélène Esnault for helpful discussions and bringing my attention to the paper [25].

(d) *Notations and conventions*

To shorten the notation, let

$$\mathbb{A} = \mathbb{A}^1 \quad \text{and} \quad \mathbb{P} = \mathbb{P}^1$$

be the affine line and the projective line, respectively in the rest of this paper. Let

$$\mathbb{D} \quad \text{and} \quad \mathbb{D}^\circ = \mathbb{D} \setminus \{0\}$$

be the open unit disc and the punctured disc of the complex plane, respectively. For a divisor D on a variety, $(D)_{\mathrm{red}}$ denotes the associated reduced subvariety, i.e., the support of D.

Since we will use the sheaves of logarithmic differentials intensively, we introduce the following notation: Let X be a smooth completion of U such that the complement $S = X \setminus U$ is a normal crossing divisor. We let

$$\breve{\Omega}^p = \breve{\Omega}^p_X = \breve{\Omega}^p_{U \subset X} := \Omega^p_X(\log S)$$

be the sheaf on X of differential forms of degree p, regular on U and with at worst logarithmic poles along S.

For a decreasing filtration F^λ indexed by $\lambda \in \mathbb{R}$, we set

$$F^{\lambda -} = \bigcap_{i<\lambda} F^i \quad \text{and} \quad F^{\lambda +} = \bigcup_{i>\lambda} F^i.$$

The λ-th graded piece $\mathrm{Gr}^\lambda = \mathrm{Gr}^\lambda_F$ of $F^\bullet$ is defined as $F^\lambda / F^{\lambda +}$.

For a complex $K = (K^\bullet, \delta^\bullet)$, the degree p term of the shift $K[n]$ is K^{n+p} with the differential δ^{n+p}. If we want to locate the degree 0 term of a complex to avoid confusion, we put the symbol ▲ under that term, e.g., $\cdots \to \underset{\blacktriangle}{A} \to B \to \cdots$. The use of ▲ in some variants in the paper should be clear. For a double complex $(K^{\bullet,\bullet}, \delta_1, \delta_2)$, the symbol $\mathrm{tot}(K^{\bullet,\bullet})$ denotes the total complex attached to $K^{\bullet,\bullet}$ with differential $\delta_1 + (-1)^p \delta_2$ on $K^{p,q}$.

When regarded as a chain map, the connection ∇ is only $\mathbb{C}$-linear and one needs to consider, for example, the derived category of sheaves of $\mathbb{C}$-vector spaces in order to discuss the cohomological properties. However when ∇ acts on $\mathcal{O}$-modules, the Godement injective resolution (of a sheaf of abelian groups) provides an injective resolution both in the category of $\mathcal{O}$-modules and of $\mathbb{C}$-vector spaces. Thus many cohomological statements can be derived within the restriction to $\mathcal{O}$-modules. On the other hand, once one reduces to consider a coherent sheaf on a projective X, one can freely work with the analytic or the Zarisky topologies by the GAGA principle. We will use both principles repeatedly without further indications.

1 The irregular Hodge filtration

(a) *The de Rham cohomology and good compactifications*

Fix a complex smooth quasi-projective variety U and a global regular function f on it, regarded as an element $f \in \mathcal{O}(U)$ or a morphism $f : U \to \mathbb{A}$ interchangeably. As in the introduction, let $\nabla = \nabla_f = d + df$ be the integrable connection on the structure sheaf $\mathcal{O}$ of U. It then extends to the twisted de Rham complex on U

$$(\Omega^\bullet, \nabla) = \Big[\mathcal{O} \xrightarrow{\nabla} \Omega^1 \xrightarrow{\nabla} \Omega^2 \to \cdots\Big].$$

Definition. The *de Rham cohomology of the connection* ∇ is the hypercohomology

$$H^i_{\mathrm{dR}}(U, \nabla) := \mathbb{H}^i\big(U, (\Omega^\bullet, \nabla)\big).$$

Difinition. Let $j : U \to X$ be a compactification of U with the complement $S := X \setminus U$. The pair (X, S) is called a *good compactification of* (U, f) if S is a normal crossing divisor of X and f extends to a morphism $f : X \to \mathbb{P}$. Thus we have the commutative diagram

$$\begin{tikzcd}
U \arrow[r, "f"] \arrow[d, "j"'] & \mathbb{A} \arrow[d] \\
X \arrow[r, "f"] & \mathbb{P}.
\end{tikzcd}$$

By the elimination of indeterminacy and the resolution of singularities there always exists a good compactification (X, S) of (U, f). Given such an X and a point $a \in f^{-1}(\infty)$ of X, there exists a system of analytically local coordinates

$$\{x_1, \cdots, x_r, t_1, \cdots, t_m, y_1, \cdots, y_l\} \quad \text{for some } r, m, l \geq 0$$

such that

- $S = (xt)$ is a union of coordinate hyperplanes, and
- $f = \frac{1}{x^e} f_0$ for some exponent $e \in \mathbb{Z}^r_{>0}$ and some analytic f_0 with $f_0(a) \neq 0$.

This local picture will be used repeatedly.

On the other hand, the connection ∇ on U extends to the twisted complex

$$(\Omega^\bullet_X(*S), \nabla) = \Big[\mathcal{O}_X(*S) \xrightarrow{\nabla} \Omega^1_X(*S) \xrightarrow{\nabla} \Omega^2_X(*S) \to \cdots\Big]. \tag{3}$$

Since $\Omega^p_X(*S) = j_*\Omega^p_U \xrightarrow{\sim} \mathcal{R}j_*\Omega^p_U$, we have

$$H^i_{\mathrm{dR}}(U, \nabla) = \mathbb{H}^i\big(X, (\Omega^\bullet_X(*S), \nabla)\big). \tag{4}$$

(*b*) *The Hodge filtration on the de Rham complex*

Fix a good compactification (X,S) of (U,f). We shall define on the complex (3) a separated filtration F^λ, indexed by $\lambda \in \mathbb{R}$, which is left continuous (i.e., $F^\lambda = F^{\lambda-}$). (It will also be exhaustive if $f: U \to \mathbb{A}$ is proper.)

Let P be the pole divisor of f on X; it is effective and supported on S. We have

$$f \in \mathcal{O}_X(P) \quad \text{and} \quad df \in \breve{\Omega}^1_X(P).$$

(Recall that $\breve{\Omega}^p_X := \Omega^p_X(\log S)$.)

Definition. Let

$$F^0(\lambda) := \Big[\mathcal{O}(\lfloor -\lambda P \rfloor) \xrightarrow{\nabla} \breve{\Omega}^1(\lfloor (1-\lambda)P \rfloor) \to \cdots \to \breve{\Omega}^p(\lfloor (p-\lambda)P \rfloor) \to \cdots\Big], \tag{5}$$

regarded as a subcomplex of (3). The *irregular Hodge filtration of* ∇ is the filtration on (3) defined by

$$F^\lambda(\nabla) = F^0(\lambda)^{\geq \lceil \lambda \rceil} \quad (\lambda \in \mathbb{R}).$$

We use $F^\lambda(\nabla)^p$ to denote the degree p component of $F^\lambda(\nabla)$; it is a locally free subsheaf of $\Omega^p_X(*S)$.

It is clear that at degree p, we have

$$F^\lambda(\nabla)^p = \begin{cases} 0 & \text{if } \lambda > p \\ \breve{\Omega}^p(\lfloor (p-\lambda)P \rfloor) & \text{if } \lambda \leq p \end{cases} \tag{6}$$

and that $F^\bullet(\nabla)$ obeys the following two rules:

$$\begin{aligned} F^\lambda(\nabla)^0 &= \left(F^{\lambda+1}(\nabla)^0\right)(P) \quad \text{if } \lambda \leq -1, \\ F^\lambda(\nabla)^p &= \breve{\Omega}^p \underset{\mathcal{O}_X}{\otimes} F^{\lambda-p}(\nabla)^0 \quad \text{for all } \lambda \in \mathbb{R}. \end{aligned} \tag{7}$$

In the rest of this subsection, we build up some basic properties of this filtration.

Proposition 1.1 *Consider the local situation. Let* $U = (\mathbb{D}^\circ)^r \times (\mathbb{D}^\circ)^m \times \mathbb{D}^l$ *with coordinates*

$$\{x_1, \cdots, x_r, t_1, \cdots, t_m, y_1, \cdots, y_l\}.$$

Let $f = \frac{f_0}{x^e} = \frac{f_0}{x_1^{e1} \cdots x_r^{er}}$ *with* $e_i > 0$, f_0 *regular and nowhere vanishing on* $\mathbb{D}^{r+m+l}$, *and* $\nabla = \nabla_f$ *the associated connection on* U. *Let* $\widetilde{U} = U \times \mathbb{D}$ *with the natural embedding into* $X = \mathbb{D}^{r+m+l+1}$ *and* $S := X \setminus \widetilde{U}$. *Then for each* λ, *the natural inclusion*

$$F^\lambda_{\boxtimes} \to F^\lambda(\widetilde{\nabla})$$

of subcomplexes of $(\Omega^\bullet_X(*S), \nabla)$ *on* X *is a quasi-isomorphism. Here* $F^\lambda_{\boxtimes}$ *is the exterior product filtration of* $F^i(\nabla)$ *and* $F^j(d)$ *while* $F^\lambda(\widetilde{\nabla})$ *is the filtration of the connection* $\widetilde{\nabla}$ *attached to* f *regarded as a function on* $\widetilde{U}$.

The same assertion remains true for $\widetilde{U} = U \times \mathbb{D}^\circ$.

Proof. Let $n = r + m + l$. Fix $\lambda \le n+1$. By a direct checking we have the natural inclusion of complexes

$$\begin{tikzcd}
F^{\lambda}_{\boxtimes}\left(\left[\mathcal{O}_{\mathbb{D}^n}(*S) \xrightarrow{\nabla} \Omega^1(*S) \to \cdots \to \Omega^n(*S)\right] \boxtimes \left[\mathcal{O}_{\mathbb{D}} \xrightarrow{d} \Omega^1\right]\right) \arrow[d] \\
F^{\lambda}\left(\mathcal{O}_X(*S) \xrightarrow{\nabla} \Omega^1(*S) \to \cdots \to \Omega^{n+1}(*S)\right).
\end{tikzcd}$$

The quotient is described as follows. Let z be the coordinate of the last piece $\mathbb{D}$ of $\widetilde{U}$. Let $A = \mathcal{O}(\widetilde{U})$ and $\Psi =$ all possible exterior products among the 1-forms in $\left\{\frac{dt_i}{t_i}, dy_j\right\}$ of degree $\ge \lambda$. Then, as an A-module, the quotient decomposes into

$$\bigoplus_{\omega\in\Psi} \left(B(\omega), \nabla|_{B(\omega)}\right)$$

with $B(\omega)[p+1] =$

$$\begin{aligned}
&\left(\frac{1}{x^{\lfloor (p+1-\lambda)e\rfloor}}A \Big/ \frac{1}{x^{\lfloor (p-\lambda)e\rfloor}}A\right)\omega_0 dz \xrightarrow{\text{left multip. by } dx^{-e}} \\
&\qquad \bigoplus_{i=1}^{r}\left(\frac{1}{x^{\lfloor (p+2-\lambda)e\rfloor}}A \Big/ \frac{1}{x^{\lfloor (p+1-\lambda)e\rfloor}}A\right)\frac{dx_i}{x_i}\omega_1 dz \\
&\qquad \to \bigoplus_{1\le i<j\le r}\left(\frac{1}{x^{\lfloor (p+3-\lambda)e\rfloor}}A \Big/ \frac{1}{x^{\lfloor (p+2-\lambda)e\rfloor}}A\right)\frac{dx_i}{x_i}\frac{dx_j}{x_j}\omega_2 dz \to \cdots
\end{aligned}$$

where $p = \deg(\omega)$ and $\omega_k = f_0^k \cdot \omega$. It is clear that the complex $B(\omega)[p+1]$ of $\mathbb{C}$-vector spaces is isomorphism to

$$\left(\frac{1}{x^{\lfloor (p+1-\lambda)e\rfloor}}A \Big/ \frac{1}{x^{\lfloor (p-\lambda)e\rfloor}}A\right) \bigotimes_{\mathbb{C}} \left[\overset{0}{\bigwedge} C \xrightarrow{\text{left multip. by } \sum_{i=1}^r v_i} \overset{1}{\bigwedge} C \to \cdots \to \overset{r}{\bigwedge} C\right]$$

where the later is the total Koszul complex attached to the $\mathbb{C}$-vector space C generated by the basis $\left\{v_i = -e_i\frac{dx_i}{x_i}\right\}_{i=1}^{r}$. Now this Koszul complex has null-cohomology and thus the assertion follows in this case.

For the case $\widetilde{U} = U \times \mathbb{D}^{\circ}$, one simply replaces dz by $\frac{dz}{z}$ in the above arguments. This then completes the proof. □

Corollary 1.2 *Consider the local situation. Let* $U = (\mathbb{D}^{\circ})^r \times (\mathbb{D}^{\circ})^m \times \mathbb{D}^l$ *with coordinates*

$$\{x_1, \cdots, x_r, t_1, \cdots, t_m, y_1, \cdots, y_l\}.$$

Let $f = \frac{f_0}{x_1^{e_1}\cdots x_r^{e_r}}$ *with* $e_i > 0$, f_0 *regular and nowhere vanishing on* $\mathbb{D}^{r+m+l}$, *and* $\nabla = \nabla_f$ *the associated connection on* U. *Let* $\widetilde{U} = U \times \mathbb{D}^a \times (\mathbb{D}^{\circ})^b$ *with the natural embedding into* $X = \mathbb{D}^{r+m+l+a+b}$ *and* $S := X \setminus \widetilde{U}$. *Then for each* λ, *the natural inclusion*

$$F^{\lambda}_{\boxtimes} \to F^{\lambda}(\widetilde{\nabla})$$

of subcomplexes of $(\Omega^\bullet_X(*S), \nabla)$ *on* X *is a quasi-isomorphism. Here* $F^\lambda_\boxtimes$ *is the exterior product filtration of* $F^i(\nabla)$ *and* $F^j(d)$ *while* $F^\lambda(\widetilde{\nabla})$ *is the filtration of the connection* $\widetilde{\nabla}$ *attached to* f *regarded as a function on* $\widetilde{U}$*.*

Proof. This follows from the above proposition and the fact that the usual Hodge filtration of the logarithmic de Rham complex of $\mathbb{D}^a \times (\mathbb{D}^\circ)^b$ is equal to the product filtration of the filtrations on its factors. □

Proposition 1.3 *Let* D *and* E *be divisors of* X *supported on* S *and* $(P)_{\text{red}}$*, respectively. Suppose* E *is effective. Then the natural inclusion*

$$\begin{tikzcd} \Big[\mathcal{O}(D) \xrightarrow{\nabla} \breve{\Omega}^1(D+P) \to \cdots \to \breve{\Omega}^p(D+pP) \to \cdots\Big] \arrow[d] \\ \Big[\mathcal{O}(D+E) \xrightarrow{\nabla} \breve{\Omega}^1(D+E+P) \to \cdots \to \breve{\Omega}^p(D+E+pP) \to \cdots\Big] \end{tikzcd} \tag{8}$$

of complexes on X *is a quasi-isomorphism.*

Proof. By induction, it suffices to consider the case where E is an irreducible component of $(P)_{\text{red}}$. The assertion is local and we may assume that $f = x^{-e} f_0$ with f_0 regular and nowhere vanishing on

$$U = (\mathbb{D}^\circ)^r \times (\mathbb{D}^\circ)^m \times \mathbb{D}^l \hookrightarrow X = \mathbb{D}^n$$

with coordinates $\{x_1, \cdots, t_1, \cdots, y_1, \cdots\}$ and $D = (x^d t^\delta), E = (x_1)$. (One has $S = (xt)$ and $(P)_{\text{red}} = (x)$.) Let $A = \mathcal{O}(\mathbb{D}^n)$ and $\Phi =$ all possible exterior products among the 1-forms in $\left\{\frac{dt_i}{t_i}, dy_j\right\}$. Then, as an A-module, the quotient complex of (8) decomposes into

$$\bigoplus_{\omega \in \Phi} \left(B(\omega), \nabla|_{B(\omega)}\right)$$

with $B(\omega)[p] =$

$$\begin{aligned} &\left(\frac{1}{x_1 x^{d+pe} t^\delta} A \Big/ \frac{1}{x^{d+pe} t^\delta} A\right) \omega_0 \xrightarrow{\text{left multip. by } dx^{-e}} \\ &\qquad \bigoplus_{i=1}^{r} \left(\frac{1}{x_1 x^{d+(p+1)e} t^\delta} A \Big/ \frac{1}{x^{d+(p+1)e} t^\delta} A\right) \frac{dx_i}{x_i} \omega_1 \\ &\qquad \to \bigoplus_{1 \le i < j \le r} \left(\frac{1}{x_1 x^{d+(p+2)e} t^\delta} A \Big/ \frac{1}{x^{d+(p+2)e} t^\delta} A\right) \frac{dx_i}{x_i} \frac{dx_j}{x_j} \omega_2 \to \cdots \end{aligned}$$

where $p = \deg(\omega)$ and $\omega_k = f_0^k \cdot \omega$. The complex $B(\omega)[p]$ of $\mathbb{C}$-vector spaces is isomorphism to

$$\left(\frac{1}{x_1 x^{d+pe} t^\delta} A \Big/ \frac{1}{x^{d+pe} t^\delta} A\right) \underset{\mathbb{C}}{\otimes} \left[\overset{0}{\bigwedge} C \xrightarrow{\text{left multip. by } \sum_{i=1}^r v_i} \overset{1}{\bigwedge} C \to \cdots \to \overset{r}{\bigwedge} C\right]$$

where the later is the total Koszul complex attached to the $\mathbb{C}$-vector space C generated by the basis $\left\{v_i = -e_i \frac{dx_i}{x_i}\right\}_{i=1}^{r}$. Now this Koszul complex has null-cohomology and hence we obtain the desired statement. $\square$

Corollary 1.4 *The inclusion*

$$\Big(F^0(\nabla) \text{ with the induced irregular Hodge filtration}\Big) \to \Big((\Omega_X^\bullet(*S), \nabla), F^\lambda(\nabla)\Big)$$

is a quasi-isomorphism of filtered complexes on X.

Proof. Write $S = (P)_{\mathrm{red}} + T$. Cor.1.2 (plus [7, proof of Prop.II.3.13]) and the above proposition give respectively the two quasi-isomorphisms

$$\begin{aligned} F^0(\nabla) &\xrightarrow{\sim} F^0(\nabla)(*T) \\ &\xrightarrow{\sim} \big(F^0(\nabla)(*T)\big)\,(*(P)_{\mathrm{red}}), \end{aligned}$$

both compatible with the equipped filtrations, while the last term is simply the complex $(\Omega_X^\bullet(*S), \nabla)$ with the filtration $F^\lambda(\nabla)$. $\square$

Notice that the corollary above implies immediately that $H^i_{\mathrm{dR}}(U, \nabla)$ is finite dimensional for any i and is non-zero only if $0 \leq i \leq 2 \cdot \dim U$ since it is the hypercohomology of a chain of coherent sheaves on a compact X of length $\dim U$.

Definition. On X the *logarithmic complex attached to* ∇ is the sub-filtered complex $(\Omega_X^\bullet(\log \nabla), F^\lambda) \subset (\Omega_X^\bullet(*S), F^\lambda(\nabla))$ defined as

$$\begin{aligned} \Omega_X^\bullet(\log \nabla) &= \Big[\ker\Big\{\mathcal{O}_X(*S) \xrightarrow{\nabla} \mathrm{Gr}^0(\nabla)^1\Big\} \xrightarrow{\nabla} F^{0+}(\nabla)\Big] \\ &= F^0(\nabla) \underset{\mathcal{O}_X}{\otimes} \mathcal{O}_X(-(P)_{\mathrm{red}}) \end{aligned}$$

where $\mathrm{Gr}^0(\nabla)^1$ is the degree 1 component of $\mathrm{Gr}^0(\nabla) = F^0(\nabla)/F^{0+}(\nabla)$.

Inside this logarithmic complex, $\Omega_X^0(\log \nabla) = \mathcal{O}(-(P)_{\mathrm{red}})$ is pure of filter degree 0 while $\Omega_X^{>0}(\log \nabla)$ is of positive filter degree, i.e. jumps > 0. The following corollary gives us the information of $\mathrm{Gr}^0(\nabla)$.

Corollary 1.5 *The inclusion* $\big(\Omega_X^\bullet(\log \nabla), \nabla, F^\lambda\big) \to \big(\Omega_X^\bullet(*S), \nabla, F^\lambda(\nabla)\big)$ *is a quasi-isomorphism of filtered complexes. In particular we have the quasi-isomorphism*

$$\mathcal{O}(-(P)_{\mathrm{red}}) \xrightarrow{\sim} \mathrm{Gr}^0(\nabla).$$

Proof. This follows from Prop.1.3 by taking $D = 0$, $E = (P)_{\mathrm{red}}$ and from the above corollary. $\square$

(*c*) *The Hodge filtration on the de Rham cohomology*

In the previous subsection, we define the irregular Hodge filtration on the twisted de Rham complex upon a chosen good compactification (X, S) of (U, f). Here we prove that the induced filtration on $H_{\mathrm{dR}}(U, \nabla)$ does not depend on the choice of X.

We begin by considering a map $\pi : (X', S') \to (X, S)$ between two good compactifications of (U, f). The corresponding irregular Hodge filtrations on them will be denoted by $F^{\lambda}_{X}(\nabla)$ and $F^{\lambda}_{X'}(\nabla)$, respectively. Recall that since π is birational, we have

$$\mathcal{R}\pi_*\mathcal{O}_{X'} = \mathcal{O}_X.$$

Lemma 1.6 *With notations as above, we have the following.*

(i) $\pi^*\breve{\Omega}^p_X = \breve{\Omega}^p_{X'}$.

(ii) *The adjuntion map* $\breve{\Omega}^p_X \to \mathcal{R}\pi_*\breve{\Omega}^p_{X'}$ *is a quasi-isomorphism.*

(iii) *We have* $\pi^*F^{\lambda}_{X}(\nabla) \subset F^{\lambda}_{X'}(\nabla)$ *where* π^* *denotes the componentwise pullback to* $\mathcal{O}_{X'}$*-modules. In particular we obtain*

$$\pi^* : \mathbb{H}\left(X, F^{\lambda}_{X}(\nabla)\right) \to \mathbb{H}\left(X', F^{\lambda}_{X'}(\nabla)\right).$$

Proof. (i) Indeed the $\mathcal{O}_{X'}$-module $\breve{\Omega}^p_{X'}$ is locally generated by forms

$$\omega = d\log f_1 \wedge \cdots \wedge d\log f_p$$

where f_j are meromorphic whose all possible zeros and poles are contained in S'. However since $X' \backslash S' = X \backslash S$, those f_j are pull-backs of meromorphic functions on X, with possible zeros and poles only on S. Thus ω is a local section of $\pi^*\breve{\Omega}^i_X$.

(ii) One has

$$\begin{aligned} \breve{\Omega}^p_X &= \mathcal{O}_X \underset{\mathcal{O}_X}{\otimes} \breve{\Omega}^p_X \\ &= \mathcal{R}\pi_*\mathcal{O}_{X'} \underset{\mathcal{O}_X}{\otimes} \breve{\Omega}^p_X \\ &\xrightarrow{\sim} \mathcal{R}\pi_*\left(\mathcal{O}_{X'} \underset{\mathcal{O}_{X'}}{\otimes} \pi^*\breve{\Omega}^p_X\right) \\ &= \mathcal{R}\pi_*\breve{\Omega}^p_{X'} \end{aligned}$$

where the isomorphism follows from the projection formula ([14, Exer.III.9.8.3] or [13, Prop.II.5.6]).

(iii) Let P and P' be the pole divisors of f on X and X', respectively. One sees readily that

$$\pi^*\mathcal{O}_X(\lfloor \eta P \rfloor) \subset \mathcal{O}_{X'}(\lfloor \eta P' \rfloor) \quad \text{for any } \eta \geq 0.$$

Thus the assertion follows from (i) and the identities in (6) and (7). □

Lemma 1.7 *Let (X, S) be a good compactification of (U, f). Suppose that $\pi : X' \to X$ is a blowup along a smooth center which has normal crossing with S. Then the canonical map*

$$F_X^\lambda(\nabla)^0 \to \mathcal{R}\pi_* \left(F_{X'}^\lambda(\nabla)^0 \right)$$

is a quasi-isomorphism for any $\lambda \in \mathbb{R}$.

Proof. Fix a non-positive real number λ. The exceptional divisor E is irreducible and is a projective bundle over the center Ξ of the blowup π. Let Q_λ be the coherent sheaf on X' defined by the short exact sequence

$$0 \to \pi^* \left(F_X^\lambda(\nabla)^0 \right) \to F_{X'}^\lambda(\nabla)^0 \to Q_\lambda \to 0. \tag{9}$$

Then Q_λ is concentrated on E. By the projection formula, it suffices to show that $\mathcal{R}\pi_* Q_\lambda$ is quasi-isomorphic to zero.

Locally on X, there exist coordinates $\{x_1, \cdots, x_n\}$ and three non-negative integers r, a, b with $a \leq r \leq b \leq n$ such that

- the function $f = (x_1^{e_1} \cdots x_r^{e_r})^{-1} f_0$ with f_0 regular and nowhere vanishing, and
- the center Ξ is defined by

$$\begin{cases} x_i = 0 & \text{if } 1 \leq i \leq a \\ x_j = 0 & \text{if } r < j \leq b. \end{cases}$$

Over this local chart, we have that

- E/Ξ is fibered by projective spaces of dimension $\dim_\Xi E = (a + b - r - 1)$,
- $\pi^* f$ has pole order $e := e_1 + \cdots + e_a$ along E, and
- above the origin of X, the sequence (9) is given by

$$0 \to \mathcal{O}\left(\lfloor -\lambda \widetilde{P} \rfloor + \sum_{i=1}^{a} \lfloor -\lambda e_i \rfloor E \right) \to \mathcal{O}\left(\lfloor -\lambda \widetilde{P} \rfloor + \lfloor -\lambda e \rfloor E \right) \to Q_\lambda \to 0 \tag{10}$$

  where $\widetilde{P} \subset X'$ denotes the proper transform of $P \subset X$. (Thus $(P')_{\text{red}} = (\widetilde{P})_{\text{red}} + E$.)

Inserting the intermediate locally free sheaves of X' into the inclusion

$$\mathcal{O}\left(\lfloor -\lambda \widetilde{P} \rfloor + \sum_{i=1}^{a} \lfloor -\lambda e_i \rfloor E \right) \subset \mathcal{O}\left(\lfloor -\lambda \widetilde{P} \rfloor + \lfloor -\lambda e \rfloor E \right)$$

by adding one more copy of the divisor E in each step, we get a filtration in the middle term of (10). It then induces a filtration on Q_λ. To get information of the induced grading on Q_λ, one has to compute the restriction

$$\mathcal{O}_E\left(\lfloor -\lambda \widetilde{P} \rfloor + \lfloor -\lambda e \rfloor E \right)$$

of the sheaf to E. Write $(\widetilde{P})_{\mathrm{red}} = \sum_{i=1}^{r} \widetilde{P}_i$ where $\widetilde{P}_i$ is the proper transform of the i-th coordinate hyperplane. We notice that, still over the origin of X,

$$E.\widetilde{P}_i = \begin{cases} H & \text{if } 1 \leq i \leq a \\ 0 & \text{if } i > a \end{cases} \quad , \quad E.E = -H \quad (H \text{ denotes a hyperplane section})$$

and

$$\begin{aligned} 0 \leq \lfloor -\lambda e \rfloor - \sum_{i=1}^{a} \lfloor -\lambda e_i \rfloor & \leq a - 1 \\ & \leq a + b - r - 1 = \dim_{\Xi} E. \end{aligned}$$

Therefore Q_λ over the origin of X is a successive extension of various $\mathcal{O}_{E/\Xi}(-\mu H)$ with $0 < \mu \leq \dim_{\Xi} E$. With μ in this range, we have that $\mathcal{R}\pi_* \mathcal{O}_{E/\Xi}(-\mu H)$ is quasi-isomorphic to zero and hence so is $\mathcal{R}\pi_* Q_\lambda$ as desired. □

Theorem 1.8 *The hypercohomology $\mathbb{H}\big(X, F^\lambda(\nabla)\big)$ only depends on (U, f), not on the choice of the good compactification (X, S).*

Proof. First suppose that $\pi : X' \to X$ is a morphism between good compactifications. Recall the weak factorization theorem of birational morphisms [28, Thm.0.0.1]: The birational morphism π admits a factorization into the following commutative diagram of birational morphisms

$$\begin{tikzcd}
X' = X_0 \arrow[r, dashed, no head] & X_1 \arrow[l, bend right] \arrow[r, dashed, no head] & \cdots \arrow[r, dashed, no head, "\alpha_m"'] & X_m \arrow[lll, bend right, "\beta_m"'] \arrow[rrr, bend left, "\gamma_m"] \arrow[r, dashed, no head] & \cdots \arrow[r, dashed, no head, "\alpha_{n-1}"'] & X_{n-1} \arrow[r, dashed, no head] \arrow[r, bend left] & X_n = X.
\arrow[from=1-1, to=1-7, bend right, "\pi"']
\end{tikzcd}$$

Here, for $1 \leq i \leq n$,

- X_i is a smooth completion of U with $S_i := X_i \setminus U$ a normal crossing divisor;
- $X_{i-1} \overset{\alpha_i}{- - -} X_i$ represents either a blowup $\alpha_i : X_{i-1} \to X_i$ of X_i along a smooth center which is of normal crossing with S_i, or a blowup $\alpha_i : X_i \to X_{i-1}$ of X_{i-1} along a smooth center which is of normal crossing with S_{i-1};
- there exists an integer $m \in [1, n]$ such that X_i are equipped with morphisms

$$\begin{aligned} \beta_i : X_i \to X_0 = X', & \quad 1 \leq i \leq m \\ \gamma_i : X_i \to X_n = X, & \quad m \leq i \leq n. \end{aligned}$$

The first and the third conditions ensure that each (X_i, S_i) is a good compactification of (U, f). We let $F_i^\lambda(\nabla)$ denote the associated irregular Hodge filtration on X_i.

Set $\gamma_0 = \pi$ and $\gamma_i = \pi \circ \beta_i : X_i \to X$ for $1 \le i \le m$. Then for each $1 \le i \le n$, we have the commutative diagrams

$$\text{(I)} \quad \begin{tikzcd} X_{i-1} \arrow[rr, "\alpha_i"] \arrow[dr, "\gamma_{i-1}"'] & & X_i \arrow[dl, "\gamma_i"] \\ & X & \end{tikzcd} \quad \text{or} \quad \text{(II)} \quad \begin{tikzcd} X_{i-1} \arrow[dr, "\gamma_{i-1}"'] & & X_i \arrow[ll, "\alpha_i"'] \arrow[dl, "\gamma_i"] \\ & X. & \end{tikzcd}$$

By Lemma 1.7 (applied to α_i), we obtain

$$\mathcal{R}\gamma_{i-1*}F^\lambda_{i-1}(\nabla)^0 = \left\{ \begin{array}{ll} \mathcal{R}\gamma_{i*}\left(\mathcal{R}\alpha_{i*}F^\lambda_{i-1}(\nabla)^0\right) & \text{in case (I)} \\ & \\ \mathcal{R}\gamma_{i-1*}\left(\mathcal{R}\alpha_{i*}F^\lambda_{i}(\nabla)^0\right) & \text{in case (II)} \end{array} \right\} = \mathcal{R}\gamma_{i*}F^\lambda_i(\nabla)^0.$$

(The $=$ means quasi-isomorphic.) Thus by induction on the index i in γ_i, one obtains that $F^\lambda_X(\nabla)^0 \to \mathcal{R}\pi_*\left(F^\lambda_{X'}(\nabla)^0\right)$ is a quasi-isomorphism. Now by the identities in (6), (7) and by Lemma 1.6(ii), we then see that $F^\lambda_X(\nabla) \to \mathcal{R}\pi_*\left(F^\lambda_{X'}(\nabla)\right)$ is a quasi-isomrphism. Therefore the assertion follows in this case.

Now given two good compactifications (X_1, S_1) and (X_2, S_2) of (U, f), one can always find a third one that dominates the two. Indeed we have the standard commutative diagram:

$$\begin{tikzcd} & X_1 & & \\ U \arrow[ur, hook] \arrow[rr, hook] \arrow[dr, hook] & & X \arrow[ul, "\pi_1"'] \arrow[dl, "\pi_2"] \arrow[r] & \overline{U} \subset X_1 \times X_2 \\ & X_2 & & \end{tikzcd}$$

where $\overline{U}$ is the closure of U in $X_1 \times X_2$ via the diagonal embedding and $X \to \overline{U}$ is a certain sequence of blowups such that $(X, X \setminus U)$ is a good compactification. The above discussion then shows that π_1 and π_2 induce isomorphisms on the hypercohomology of the corresponding $F^\lambda(\nabla)$. This completes the proof. $\square$

Applying the snake lemma to the long exact sequence associated with

$$0 \to F^{\lambda+}(\nabla) \to F^\lambda(\nabla) \to \mathrm{Gr}^\lambda(\nabla) \to 0,$$

the above theorem then yields the following.

Corollary 1.9 *The hypercohomology* $\mathbb{H}^i\left(X, \mathrm{Gr}^\lambda(\nabla)\right)$ *does not depend on the choice of* X.

Definition. Let (X, S) be a good compactification of (U, f).

(i) For any $\lambda \in \mathbb{R}$, we define

$$\begin{aligned} H^i\Big(U, F^\lambda(\nabla)\Big) &:= \mathbb{H}^i\Big(X, F^\lambda(\nabla)\Big) \\ H^i\Big(U, \mathrm{Gr}^\lambda(\nabla)\Big) &:= \mathbb{H}^i\Big(X, \mathrm{Gr}^\lambda(\nabla)\Big). \end{aligned}$$

(ii) The *irregular Hodge filtration* F^λ on $H^i_{\mathrm{dR}}(U,\nabla)$ is defined by setting

$$\begin{aligned} F^\lambda H^i_{\mathrm{dR}}(U,\nabla_f) &= \text{Image}\left\{\mathbb{H}^i\left(X, F^\lambda(\nabla)\right) \to \mathbb{H}^i\left(X, (\Omega^\bullet_X(*S),\nabla)\right)\right\} \\ &= \text{Image}\left\{H^i\left(U, F^\lambda(\nabla)\right) \to H^i_{\mathrm{dR}}(U,\nabla)\right\} \end{aligned}$$

induced from the inclusion $F^\lambda(\nabla) \to (\Omega^\bullet_X(*S),\nabla)$ and via the canonical isomorphism (4).

The definition does not depend on the choice of X. Notice that by Cor.1.4 we have

$$H^i\left(U, F^\lambda(\nabla)\right) = H^i_{\mathrm{dR}}(U,\nabla) \quad \text{if } \lambda \le 0.$$

(*d*) *The cohomology with compact support*

In the last part of this section we introduce the de Rham cohomology with compact support of the connection ∇ and define the corresponding irregular Hodge filtration. For the classical case, see [10, §4.3]. Again the definitions rely on choosing a good compactification X first. It is possible to establish the corresponding properties for the cohomology with compact support and prove that the definition of the irregular Hodge filtration does not depend on the choice of X as in the previous discussion. However we do not proceed in this direction. The independency will be clear once we obtain the duality in the next section. Notice that the proofs of the results in the next section do not use the proposition below.

Definition. Let (X,S) be a good compactification of (U,f) and P be the pole divisor of f on X.

(i) The *de Rham cohomology of* (U,∇) *with compact support* is the hypercohomology

$$H^i_{\mathrm{dR,c}}(U,\nabla) = \mathbb{H}^i\left(X, \mathcal{O}(-S) \xrightarrow{\nabla} \breve{\Omega}^1(-S+P) \to \cdots \to \breve{\Omega}^p(-S+pP) \to \cdots\right).$$

(ii) Write $S = (P)_{\mathrm{red}} + T$. For any $\lambda \in \mathbb{R}$, define

$$F^\lambda_{\mathrm{c}}(\nabla) := \left(F^\lambda(\nabla)\right)(-T),$$

regarded as a subcomplex of $F^\lambda(\nabla)$ on X. (The stability under ∇ in $F^\lambda_{\mathrm{c}}(\nabla)$ is easy to check.) Set $\mathrm{Gr}^\lambda_{\mathrm{c}}(\nabla) = F^\lambda_{\mathrm{c}}(\nabla)/F^{\lambda+}_{\mathrm{c}}(\nabla)$. We let

$$\begin{aligned} H^i_{\mathrm{c}}\left(U, F^\lambda(\nabla)\right) &:= \mathbb{H}^i\left(X, F^\lambda_{\mathrm{c}}(\nabla)\right) \\ H^i_{\mathrm{c}}\left(U, \mathrm{Gr}^\lambda(\nabla)\right) &:= \mathbb{H}^i\left(X, \mathrm{Gr}^\lambda_{\mathrm{c}}(\nabla)\right). \end{aligned}$$

(iii) By Prop.1.3 (for $D = 0, E = (P)_{\mathrm{red}}$), we have

$$H^i_{\mathrm{dR,c}}(U, \nabla) = H^i_{\mathrm{c}}\left(U, F^0(\nabla)\right).$$

The *irregular Hodge filtration* on $H^i_{\mathrm{dR,c}}(U, \nabla)$ is the filtration

$$\begin{aligned} F^\lambda H^i_{\mathrm{dR,c}}(U, \nabla) &= \mathrm{Image}\left\{\mathbb{H}^i\left(X, F^\lambda_{\mathrm{c}}(\nabla)\right) \to \mathbb{H}^i\left(X, F^0_{\mathrm{c}}(\nabla)\right)\right\} \\ &= \mathrm{Image}\left\{H^i_{\mathrm{c}}\left(U, F^\lambda(\nabla)\right) \to H^i_{\mathrm{dR,c}}(U, \nabla)\right\}. \end{aligned}$$

In particular, if $f : U \to \mathbb{A}$ is proper (i.e., $S = (P)_{\mathrm{red}}$), one has the natural isomorphism

$$\left(H^i_{\mathrm{dR,c}}(U, \nabla), F^\lambda\right) \xrightarrow{\sim} \left(H^i_{\mathrm{dR}}(U, \nabla), F^\lambda\right).$$

Proposition 1.10 *Let U, f, ∇ be as before. We have the following functorial properties.*

(i) *Let $a : U' \to U$ be a proper morphism of smooth quasi-projective varieties and let $\nabla' = a^*\nabla$ be the pullback connection on U'. Then the natural map $a^*(\Omega^\bullet_U, \nabla) \to (\Omega^\bullet_{U'}, \nabla')$ induces*

$$a^* : H^q_{\mathrm{c}}\left(U, F^\lambda(\nabla)\right) \to H^q_{\mathrm{c}}\left(U', F^\lambda(\nabla')\right).$$

(ii) *Let $i : V \to U$ be a smooth divisor and $j : U^\circ \to U$ be the complement. Then we have the natural long exact sequence*

$$\cdots \to H^q_{\mathrm{c}}\left(U^\circ, F^\lambda(\nabla)\right) \xrightarrow{j_*} H^q_{\mathrm{c}}\left(U, F^\lambda(\nabla)\right) \xrightarrow{i^*} H^q_{\mathrm{c}}\left(V, F^\lambda(\nabla)\right) \to \cdots.$$

Proof. (i) Take a compactification $b : X' \to X$ of $a : U' \to U$ such that $(X', X' \setminus U')$ and $(X, X \setminus U)$ are good compactifications of $(U', f \circ a)$ and (U, f), respectively. Write $X' \setminus U' = T' + (P')_{\mathrm{red}}$ where P' is the pole divisor of $f \circ a$. Then we have $T' \subset b^{-1}(T)$ since a is proper. Thus $b^* F^\lambda_{\mathrm{c}}(\nabla) \subset F^\lambda_{\mathrm{c}}(\nabla')$ and the assertion follows.

(ii) Choose a good compactification (X, S) of (U, f) such that $S + \bar{V}$ forms a normal crossing divisor of X where $\bar{V}$ is the closure of $V \subset X$. Thus $(X, S + \bar{V})$ and $(\bar{V}, S \cap \bar{V})$ are good for (U°, f) and (V, f), respectively. On X, we have

$$F^\lambda_{\mathrm{c}}(\nabla|_{U^\circ}) = F^\lambda_{\mathrm{c}}(\nabla)(-\bar{V}) \subset F^\lambda_{\mathrm{c}}(\nabla).$$

Moreover the natural sequence

$$0 \to F^\lambda_{\mathrm{c}}(\nabla|_{U^\circ}) \to F^\lambda_{\mathrm{c}}(\nabla) \xrightarrow{i^*} F^\lambda_{\mathrm{c}}(\nabla|_V) \to 0$$

is exact as can be derived easily by local computations. (Cf. [24, Example 7.23(1)] and [10, Prop.3.7.15] for the case f is trivial. In both references, V is allowed to be a normal crossing divisor.) The assertion then follows by taking hypercohomology. □

2 The duality

In this section we assume that U is irreducible of $\dim U = n$. We shall define canonically a perfect linear pairing

$$H^i_{\mathrm{dR}}(U, \nabla_f) \times H^{2n-i}_{\mathrm{dR,c}}(U, \nabla_{-f}) \xrightarrow{\langle\!\langle\, ,\rangle\!\rangle} H^{2n}_{\mathrm{dR,c}}(U) = \mathbb{C}$$

for every i, which is compatible with the irregular Hodge filtrations on them.

(a) *The pairing on the de Rham cohomology*

To define the pairing on the de Rham cohomology of (U, ∇), we mimic Deligne's construction in [9, p.124].

Fix a complex smooth irreducible quasi-projective variety U of dimension n and a regular function $f \in \mathcal{O}(U)$. Let (X, S) be a fixed good compactification of (U, f) throughout the discussion. As before, let P be the pole divisor of f on X and write

$$S = (P)_{\mathrm{red}} + T.$$

We construct a chain map

$$F^0(\nabla_f) \underset{\mathcal{O}_X}{\otimes} \big(F^0_{\mathrm{c}}(\nabla_{-f})(-(P)_{\mathrm{red}})\big) \xrightarrow{\langle\, ,\rangle} (\Omega^\bullet_X, d) \tag{11}$$

to the usual de Rham complex of X in the following way. First by Prop.1.3, the inclusions of complexes

$$F^0(n) = F^0(\nabla_f)(-nP) \hookrightarrow F^0(\nabla_f)$$

is a quasi-isomorphism and we have a chain map from $F^0(n) \otimes F^0(\nabla_{-f})(-(P)_{\mathrm{red}})$:

$$\begin{tikzcd}
\mathcal{O}(-nP) \xrightarrow{\nabla_f} \cdots \to \breve{\Omega}^{n-i}(-iP) \to \cdots \to \breve{\Omega}^n \\
\otimes \\
\mathcal{O}(-S) \xrightarrow{\nabla_{-f}} \cdots \to \breve{\Omega}^j(-S+jP) \to \cdots \to \breve{\Omega}^n(-S+nP) \arrow[d] \\
\mathcal{O}(-S-nP) \xrightarrow{d} \cdots \to \breve{\Omega}^{n-k}(-S-kP) \to \cdots \to \breve{\Omega}^n(-S) = \Omega^n.
\end{tikzcd} \tag{12}$$

Here the pairings

$$\breve{\Omega}^{n-i}(-iP) \underset{\mathcal{O}_X}{\otimes} \breve{\Omega}^j(-S+jP) \to \breve{\Omega}^{n-i+j}(-S-(i-j)P)$$

appeared in the above chain map are the natural exterior product. Now the last complex in (12) is a subcomplex of $(\Omega^\bullet_X, d)$. Thus, via this inclusion, we obtain the desired chain map (11).

Taking hypercohomology, we then obtain the Poincaré pairing

$$\begin{array}{rcl} H^i_{\mathrm{dR}}(U,\nabla)\times H^{2n-i}_{\mathrm{dR,c}}(U,\nabla_{-f}) \xrightarrow{\langle\!\langle\,,\rangle\!\rangle} H^{2n}_{\mathrm{dR}}(X)=H^n(X,\Omega^n) & \xrightarrow{\sim} & \mathbb{C} \\ \text{continuous } (n,n)\text{-form } \omega & \mapsto & \displaystyle\int_X \omega. \end{array}$$

Theorem 2.1 *For any i, the Poincaré pairing $\langle\!\langle\,,\rangle\!\rangle$ constructed above is perfect.*

Proof. Indeed we have the perfect pairing

$$\breve{\Omega}^i(mS)\otimes\breve{\Omega}^{n-i}(m'S)\to\breve{\Omega}^n((m+m')S)=\Omega^n_X((m+m'+1)S).$$

Consider the Hom-sheaf with value in Ω^n_X

$$(\bullet)^\wedge := \underline{\mathrm{Hom}}_{\mathcal{O}_X}(\bullet,\Omega^n_X).$$

Then we have

$$\begin{aligned} \left(F^0_{\mathrm{c}}(\nabla_{-f})\right)^\wedge &= \left[\mathcal{O}\xrightarrow{\nabla_f}\cdots\to \underset{\blacktriangle}{\breve{\Omega}^n(nP)}\right]\otimes\mathcal{O}(-nP) \\ &\simeq F^0(\nabla_f)[n] \quad \text{(Prop.1.3)}. \end{aligned}$$

Therefore by filtering the complexes and the Serre duality, we have

$$H^i_{\mathrm{dR,c}}(U,\nabla_{-f})^\vee=\mathbb{H}^{n-i}\left(X,F^0(\nabla_f)[n]\right)=H^{2n-i}_{\mathrm{dR}}(U,\nabla_f).$$

Here $(\bullet)^\vee$ denotes the dual vector space.

One can use, e.g., the fine resolution of the twisted de Rham complex into sheaves of continuous (p,q)-forms (with appropriate poles along S) to check that the argument here is compatible with the definition of the pairing. (Cf. the proof of the next theorem.) □

Theorem 2.2 *The two pairs $(H^i_{\mathrm{dR}}(U,\nabla_f),F^\bullet)$ and $(H^{2n-i}_{\mathrm{dR,c}}(U,\nabla_{-f}),F^\bullet)$ of filtered vector spaces are dual to each other via the perfect Poincaré pairing (up to a degree shift). More precisely, for any λ we have*

$$\langle\!\langle F^\lambda H^i_{\mathrm{dR}}(\nabla_f),F^{(n-\lambda)+}H^{2n-i}_{\mathrm{dR,c}}(\nabla_{-f})\rangle\!\rangle=0=\langle\!\langle F^{\lambda+}H^i_{\mathrm{dR}}(\nabla_f),F^{(n-\lambda)}H^{2n-i}_{\mathrm{dR,c}}(\nabla_{-f})\rangle\!\rangle \tag{13}$$

and the Poincaré pairing induces a duality between $\mathrm{Gr}^\lambda H^i_{\mathrm{dR}}(\nabla_f)$ and $\mathrm{Gr}^{n-\lambda} H^{2n-i}_{\mathrm{dR,c}}(\nabla_{-f})$. (We have omitted the base U inside the cohomology in the formulas.)

Proof. We use the fine resolution into continuous (p,q)-forms. Suppose $\omega\in F^\lambda H^i_{\mathrm{dR}}(\nabla_f)$. Since ω is the image of an element in $H^i(F^\lambda(\nabla_f))$, it is represented by

$$\sum\omega_p\in\bigoplus_{p\geq\lambda}\Gamma\left(X,\breve{\Omega}^{p,i-p}_\infty(\lfloor(p-\lambda)P\rfloor)\right)\subset\bigoplus_{p\geq 0}\Gamma\left(X,\breve{\Omega}^{p,i-p}_\infty(\lfloor(p-\lambda)P\rfloor)\right).$$

Here $\breve{\Omega}^{p,i-p}_\infty$ denotes the sheaf of logarithmic $(p,i-p)$-forms with infinitely differentiable coefficients. As the inclution

$$\left(\breve{\Omega}^p(-(P)_{\mathrm{red}}+\lfloor(p-\lambda)P\rfloor),\nabla_f\right)_{p\geq0}\subset\left(\breve{\Omega}^p(\lfloor(p-\lambda)P\rfloor),\nabla_f\right)_{p\geq0}$$

is a quasi-isomorphism, there exists $\alpha \in \bigoplus_{p\geq 0} \Gamma\left(X, \breve{\Omega}_\infty^{p,i-1-p}(\lfloor (p-\lambda)P\rfloor)\right)$ such that

$$\sum(\omega + D(\alpha))_p \in \bigoplus_{p\geq 0} \Gamma\left(X, \breve{\Omega}_\infty^{p,i-p}(-(P)_{\mathrm{red}} + \lfloor (p-\lambda)P\rfloor)\right).$$

Here D is the total differential given by

$$D = \nabla + (-1)^p \bar{d} \quad \text{on } \breve{\Omega}_\infty^{pq}(rP).$$

Now given $\eta \in F^{(n-\lambda)+} H_c^{2n-i}(\nabla_{-f})$, which is represented as the sum

$$\sum \eta_q \in \bigoplus_{q>n-\lambda} \Gamma\left(X, \breve{\Omega}_\infty^{q,2n-i-q}(-T + \lfloor (q-n+\lambda-\varepsilon)P\rfloor)\right) \quad \text{for some } \varepsilon > 0,$$

one obtains

$$\begin{aligned} \langle\!\langle \omega, \eta \rangle\!\rangle &= \sum_{q>n-\lambda} \langle\!\langle (\omega + D(\alpha))_{n-q}, \eta_q \rangle\!\rangle \quad \text{(definition)} \\ &= \sum_{q>n-\lambda} \langle\!\langle D(\alpha)_{n-q}, \eta_q \rangle\!\rangle \quad (\text{as } \omega_{n-q} = 0 \text{ if } n-q < \lambda) \\ &= \int_X D(\alpha \wedge \eta) \quad (\text{as } D\eta = 0) \\ &= 0 \quad \text{(Stokes)}. \end{aligned}$$

The application of the Stokes theorem in the last equality above is valid since the continuous $(2n-1)$-form $\alpha \wedge \eta$ has no poles on the compact X. Indeed for $q > n - \lambda$ and any $\varepsilon > 0$,

$$\begin{aligned} \alpha_{n-q} \wedge \eta_q &\in \Gamma\left(X, \breve{\Omega}_\infty^{n,n-1}(\lfloor (n-q-\lambda)P\rfloor - T + \lfloor (q-n+\lambda-\varepsilon)P\rfloor)\right) \\ &\subset \Gamma\left(X, \breve{\Omega}_\infty^{n,n-1}(-(P)_{\mathrm{red}} - T)\right) \\ &= \Gamma\left(X, \Omega_\infty^{n,n-1}\right), \end{aligned}$$

while similarly

$$\begin{aligned} \alpha_{n-1-q} \wedge \eta_q &\in \Gamma\left(X, \breve{\Omega}_\infty^{n-1,n}(\lfloor (n-1-q-\lambda)P\rfloor - T + \lfloor (q-n+\lambda-\varepsilon)P\rfloor)\right) \\ &\subset \Gamma\left(X, \breve{\Omega}_\infty^{n-1,n}(-(P)_{\mathrm{red}} - T)\right) \\ &\subset \Gamma\left(X, \Omega_\infty^{n-1,n}\right). \end{aligned}$$

The second equality of (13) can be proved similarly. □

Corollary 2.3 *The filtered cohomology* $(H^i_{\mathrm{dR,c}}(U, \nabla), F^\lambda)$ *with compact support does not dependent on the choice of the compactification* X.

(b) *Pairings on the spectral sequence*

Again we have $S := X \setminus U$ containing the support $(P)_{\mathrm{red}}$ of the poles of f and $S = (P)_{\mathrm{red}} + T$. Take a sequence

$$\lambda_{-1} < 0 = \lambda_0 < \lambda_1 < \cdots < \lambda_N = n < \lambda_{N+1} \quad \text{with } \lambda_i + \lambda_{N-i} = n$$

where $\lambda_0, \cdots, \lambda_N$ are all the non-negative jumps of the filtration $F^\lambda(\nabla)$ on X. Notice that we have

$$\lfloor \lambda_{i-1} P \rfloor + \lfloor -\lambda_i P \rfloor = -(P)_{\mathrm{red}}$$

as can be checked easily. The associated Hodge to de Rham spectral sequence reads

$$E_{1,*}^{p,q} = H_*^{p+q}\left(U, \mathrm{Gr}^{\lambda_p}(\nabla)\right) \Longrightarrow H^{p+q}_{\mathrm{dR},*}(U, \nabla)$$

where $* = \mathrm{c}$ or nothing.

E_r-terms and jumping gradings

For $0 \le i < j \le N+1$, let

$$G\binom{i}{j} := \mathrm{tot}\begin{bmatrix} F^{\lambda_j}(\nabla_f) \\ \downarrow \\ \underset{\blacktriangle}{F^{\lambda_i}(\nabla_f)} \end{bmatrix} \quad \text{and} \quad G_{\mathrm{c}}\binom{i}{j} := \mathrm{tot}\begin{bmatrix} F_{\mathrm{c}}^{\lambda_j}(\nabla_{-f}) \\ \downarrow \\ \underset{\blacktriangle}{F_{\mathrm{c}}^{\lambda_i}(\nabla_{-f})} \end{bmatrix}$$

be the *jumping gradings*, which are the representatives of the quotients $F^{\lambda_i}(\nabla_f)/F^{\lambda_j}(\nabla_f)$ and $F_{\mathrm{c}}^{\lambda_i}(\nabla_{-f})/F_{\mathrm{c}}^{\lambda_j}(\nabla_{-f})$, respectively. Notice that similar to Cor.1.9 the hypercohomology of $G\binom{i}{j}$ over X is independent of the choice of X.

For two pairs of numbers $i < j$ and $i' < j'$, we say $(i, j) \ge (i', j')$ if $i \ge i'$ and $j \ge j'$. Then for any $(i, j) \ge (i', j')$, there is the natural componentwise inclusion

$$G\binom{i}{j} \to G\binom{i'}{j'} \tag{14}$$

and if $i < j < k$, one has the distinguished triangle

$$G\binom{j}{k} \to G\binom{i}{k} \to G\binom{i}{j} \xrightarrow{+1} .$$

We have

$$E_r^{p,q-p} = \mathrm{Image}\left\{\mathbb{H}^q\left(X, G\binom{p}{p+r}\right) \to \mathbb{H}^q\left(X, G\binom{p-r+1}{p+1}\right)\right\},$$

traditionally regarded as a subquotient of $\mathbb{H}^q\left(X, G\binom{p}{p+1}\right)$. The following commutative diagram illustrates the various terms in the spectral sequence.

$$\begin{tikzcd}
\mathbb{H}^q(G\binom{p}{N+1}) \arrow[r] \arrow[dddrrrr, bend right, "\text{Image} = E_\infty"'] & \mathbb{H}^q(G\binom{p}{N}) \arrow[r] & \cdots \arrow[r] & \mathbb{H}^q(G\binom{p}{p+2}) \arrow[r] \arrow[dr, bend right, "\text{Image} = E_2"'] & \mathbb{H}^q(G\binom{p}{p+1}) = E_1^{p,q-p} \arrow[d] \\
 & & & & \mathbb{H}^q(G\binom{p-1}{p+1}) \arrow[d] \\
 & & & & \vdots \arrow[d] \\
 & & & & \mathbb{H}^q(G\binom{0}{p+1})
\end{tikzcd} \tag{15}$$

(Here we omit the base X of the hypercohomology. Notice that $G\binom{p}{N+1} = F^{\lambda_p}(\nabla_f)$ and $G\binom{p}{p+1} = \mathrm{Gr}^{\lambda_p}(\nabla_f)$.)

Similar pictures hold for $G_{\mathrm{c}}\binom{i}{j}$ (but replace ∇_f by ∇_{-f} in this case).

From subs to quots

Recall the complex $F^0(\lambda)$ defined in (5) of §1. Notice that by Prop.1.3 the inclusion

$$F^0(\lambda) = F^0(\nabla)(\lfloor -\lambda P \rfloor) \to F^0(\nabla)$$

is a quasi-isomorphism for any $\lambda \geq 0$. Define

$$Q_\lambda = F^0(\lambda)^{<\lceil \lambda \rceil}.$$

We have the short exact sequence

$$0 \to F^\lambda(\nabla_f) \to F^0(\lambda) \to Q_\lambda \to 0.$$

Now for $0 \leq i < j \leq N+1$, define

$$Q\begin{pmatrix} i \\ j \end{pmatrix} := \operatorname{tot}\begin{bmatrix} Q_{\lambda_j} \\ \downarrow \\ \underset{\blacktriangle}{Q_{\lambda_i}} \end{bmatrix}.$$

Then for $(i,j) \geq (i',j')$, we have the componentwise quotient

$$Q\begin{pmatrix} i \\ j \end{pmatrix} \to Q\begin{pmatrix} i' \\ j' \end{pmatrix} \tag{16}$$

and the distinguished triangles

$$G\begin{pmatrix} i \\ j \end{pmatrix} \to \operatorname{tot}\begin{bmatrix} F^0(\lambda_j) \\ \downarrow \\ \underset{\blacktriangle}{F^0(\lambda_i)} \end{bmatrix} \to Q\begin{pmatrix} i \\ j \end{pmatrix} \xrightarrow{+1} .$$

Since the middle term is quasi-isomorphic to zero, we obtain a quasi-isomorphism

$$Q\binom{i}{j}[-1] \xrightarrow{\sim} G\binom{i}{j}. \tag{17}$$

The natural maps (14) and (16) are compatible with the above quasi-isomorphism.

The pairings

To show that the spectral sequence is compatible with the duality, we should construct pairings

$$\langle\,,\rangle^i_j : G\binom{i}{j} \underset{\mathcal{O}_X}{\otimes} G_{\mathrm{c}}\binom{N+1-j}{N+1-i} \to (\Omega^\bullet_X, d) \tag{18}$$

for all $0 \le i < j \le N+1$ which induce perfect pairings on cohomology and are compatible with respect to the partial ordering of various (i,j).

Define a pairing

$$Q\binom{i}{j}[-1] \underset{\mathcal{O}_X}{\otimes} G_{\mathrm{c}}\binom{N+1-j}{N+1-i} \xrightarrow{\langle\,,\rangle^i_j} (\Omega^\bullet_X, d)$$

as follows. For ω in the degree p term of $Q\binom{i}{j}[-1]$

$$\omega = (\omega_1, \omega_2) \in \breve{\Omega}^p\big(\lfloor (p-\lambda_j)P\rfloor\big) \oplus \breve{\Omega}^{p-1}\big((p-1-\lambda_i)P\big) \tag{19}$$

and η in the degree q term of $G_{\mathrm{c}}\binom{N+1-j}{N+1-i}$

$$\begin{aligned} \eta &= (\eta_1, \eta_2) \\ &\in \breve{\Omega}^{q+1}\big(-T + \lfloor (q+1-\lambda_{N+1-i})P\rfloor\big) \oplus \breve{\Omega}^q\big(-T + \lfloor (q-\lambda_{N+1-j})P\rfloor\big) \\ &= \breve{\Omega}^{q+1}\big(-T + \lfloor (q+1-n+\lambda_{i-1})P\rfloor\big) \oplus \breve{\Omega}^q\big(-T + \lfloor (q-n+\lambda_{j-1})P\rfloor\big), \end{aligned} \tag{20}$$

we set

$$\begin{aligned} \langle \omega, \eta\rangle^i_j := \omega_1\eta_2 + (-1)^q\omega_2\eta_1 \quad &\in \quad \breve{\Omega}^{p+q}\big(-T + \lfloor \lambda_{i-1}P\rfloor + \lfloor -\lambda_i P\rfloor + (p+q-n)P\big) \\ &= \quad \breve{\Omega}^{p+q}\big(-S - (n-p-q)P\big) \\ &\subset \quad \Omega^{p+q}_X. \end{aligned}$$

One checks readily that

$$d\langle \omega, \eta\rangle^i_j = \langle \nabla_f(\omega), \eta\rangle^i_j + (-1)^p \langle \omega, \nabla_{-f}(\eta)\rangle^i_j,$$

where we have adapted the sign convention

$$\begin{aligned} \nabla_f(\omega_1, \omega_2) &= \big(\nabla_f(\omega_1), (-1)^{p-1}\omega_1 + \nabla_f(\omega_2)\big) \\ \nabla_{-f}(\eta_1, \eta_2) &= \left(\nabla_{-f}(\eta_1), (-1)^q\eta_1 + \nabla_{-f}(\eta_2)\right). \end{aligned}$$

(The sign $(-1)^{p-1}$ in the first equation is due to the shift $[-1]$ in $Q\binom{i}{j}[-1]$.)

Using the quasi-isomorphism (17) we obtain the pairing (18) which induces a pairing

$$\mathbb{H}^q\left(X, G\binom{i}{j}\right) \times \mathbb{H}^{2n-q}\left(X, G_{\mathrm{c}}\binom{N+1-j}{N+1-i}\right) \xrightarrow{\langle\!\langle\,,\rangle\!\rangle^i_j} H^{2n}_{\mathrm{dR}}(X) = \mathbb{C} \tag{21}$$

for each q.

Theorem 2.4 *For all* $0 \leq i < j \leq N+1$*, the parings* $\langle\!\langle\, , \rangle\!\rangle^i_j$ *are perfect and they are compatible with each other under the partial ordering of* (i,j) *and the map* (14).

Proof. The proof of the perfectness is similar to that of Thm.2.1. One shows that the cohomology of the various truncations

$$Q\binom{i}{j}[-1]^{\geq n-l} \quad \text{and} \quad G_{\mathrm{c}}\binom{N+1-j}{N+1-i}^{\leq l}$$

are dual to each other via the pairing by induction on the length l. In each step, the perfectness follows from the classical Serre duality asserting the perfectness of the pairing

$$H^q\left(X, \breve{\Omega}^p(D)\right) \times H^{n-q}\left(X, \breve{\Omega}^{n-p}(-S-D)\right) \to H^n(X, \Omega^n) = H^{2n}_{\mathrm{dR}}(X).$$

The compatibilities with the ordering of (i,j) and with the definition of the parings are clear by e.g. writing everything in terms of continuous differential forms. □

Remark. For $(i,j) = (0, N+1)$, the quasi-isomorphism (17) reduces to the inclusion $F^0(\lambda_{N+1}) \to F^0(\nabla)$. One can use this inclusion and the complex $F^0_{\mathrm{c}}(\nabla_{-f})$ instead of $F^0(n) \to F^0(\nabla)$ and $F^0_{\mathrm{c}}(\nabla_{-f})(-(P)_{\mathrm{red}})$, respectively in the construction (12) of the previous subsection to define the (same) pairing on the de Rham cohomology. In this case the above theorem then recovers Thm.2.1.

Corollary 2.5 *The cohomology* $H^q_{\mathrm{c}}\left(U, F^\lambda(\nabla_f)\right)$ *and* $H^q_{\mathrm{c}}\left(U, \mathrm{Gr}^\lambda(\nabla_f)\right)$ *do not depend on the choice of the good compactification* (X,S) *of* (U,f)*.*

Proof. Applying the above theorem for $i=0$, we see that $H^{2n-q}_{\mathrm{c}}\left(U, F^{\lambda_{N+1-j}}(\nabla_{-f})\right)$ is canonically dual to $\mathbb{H}^q\left(X, G\binom{0}{j}\right)$. As already mentioned, this later space is independent of the choice of X. Thus after renaming the indices and the function f, we see that $H^q_{\mathrm{c}}\left(U, F^\lambda(\nabla_f)\right)$ is independent of X. The other statement follows by taking long exact sequence and from the compatibility of the pairings with the ordering of (i,j). □

Using the description in (15), the above theorem and the remark after it imply the following.

Corollary 2.6 *The pairings* $\langle\!\langle\, , \rangle\!\rangle^i_j$ *induce perfect pairings*

$$\langle\!\langle\, , \rangle\!\rangle_r : E^{p,q}_r \times E^{N-p, 2n-N-q}_{r,\mathrm{c}} \to \mathbb{C}.$$

They are compatible with the pairing on the de Rham cohomology in the sense that

$$\langle\!\langle \omega, \eta \rangle\!\rangle_r = \langle\!\langle \omega, \eta \rangle\!\rangle$$

for $(\omega, \eta) \in H^{p+q}\left(U, F^{\lambda_p}(\nabla)\right) \times H^{2n-p-q}_{\mathrm{c}}\left(U, F^{n-\lambda_p}(\nabla)\right)$ *projected into* E_r*-term in the left and into the de Rham cohomology in the right.*

Proposition 2.7 *For $0 \le i < j < k \le N+1$, the sequence of pairings constructed in* (21) *on the cohomology of the two distinguished triangles*

$$\begin{tikzcd}
G\binom{j}{k} \arrow[r] & G\binom{i}{k} \arrow[r] & G\binom{i}{j} \arrow[r, "+1"] & {}
\end{tikzcd}$$

$$\begin{tikzcd}
{} & G_{\mathrm{c}}\binom{N+1-k}{N+1-j} \arrow[l, "+1"'] & G_{\mathrm{c}}\binom{N+1-k}{N+1-i} \arrow[l] & G_{\mathrm{c}}\binom{N+1-j}{N+1-i} \arrow[l]
\end{tikzcd}$$

commutes up to sign.

Proof. We show that already in the chain level the diagram

$$\begin{tikzcd}
G\binom{i}{j} \arrow[d, "\delta"'] & \otimes\, G_{\mathrm{c}}\binom{N+1-j}{N+1-i} \arrow[r, "\langle\,,\rangle^i_j"] & (\Omega^\bullet_X, d) \arrow[d, equal] \\
G\binom{j}{k} & \otimes\, G_{\mathrm{c}}\binom{N+1-k}{N+1-j} \arrow[u, "\delta_{\mathrm{c}}"'] \arrow[r, "\langle\,,\rangle^j_k"] & (\Omega^\bullet_X, d)
\end{tikzcd}$$

commutes up to sign where δ and δ_{c} are the map $+1$ in the distinguished triangles. Let ω and η be degree p and q elements in $Q\binom{i}{j}[-1]$ and $G_{\mathrm{c}}\binom{N+1-k}{N+1-j}$ as in (19) and (20), respectively (but replace (i,j) by (j,k) in (20)). Then

$$\begin{aligned}
\delta(\omega) &= (0, \omega_1) \\
&\qquad \in \breve{\Omega}^{p+1}\big(\lfloor (p+1-\lambda_k)P \rfloor\big) \oplus \breve{\Omega}^{p}\big((p-\lambda_j)P\big), \\
\delta_{\mathrm{c}}(\eta) &= (0, \eta_1) \\
&\qquad \in \breve{\Omega}^{q+2}\big(-T + \lfloor (q+2-n+\lambda_{i-1})P \rfloor\big) \oplus \breve{\Omega}^{q+1}\big(-T + \lfloor (q+1-n+\lambda_{j-1})P \rfloor\big).
\end{aligned}$$

Thus

$$\langle \delta(\omega), \eta \rangle^j_k = (-1)^q \omega_1 \eta_1 = (-1)^q \langle \omega, \delta_{\mathrm{c}}(\eta) \rangle^i_j.$$

On the cohomology level, we then have

$$\langle\!\langle \delta(\omega), \eta \rangle\!\rangle^j_k + (-1)^p \langle\!\langle \omega, \delta_{\mathrm{c}}(\eta) \rangle\!\rangle^i_j = 0$$

for $\omega \in \mathbb{H}^p\left(X, G\binom{i}{j}\right)$ and $\eta \in \mathbb{H}^{2n-p-1}\left(X, \binom{N+1-k}{N+1-j}\right)$. ☐

Remark. The above proposition reduces to the special case of the compatibility of the pairings with respect to the ordering of (i,j) in the previous theorem if the spectral sequence degenerates at the E_1-terms since the connection maps in cohomology induced by δ and δ_{c} in the proof are then all zero.

3 A surface example

In this section, we consider the following situation. Let X be a complex smooth projective (irreducible) surface, which is *regular*, i.e.,

$$H^1(X, \mathcal{O}_X) = 0.$$

Let $f : X \to \mathbb{P}$ be a morphism such that the fiber $S := f^{-1}(\infty)$ is a connected (reduced) normal crossing divisor on X. Fix a positive integer e. We consider the irregular Hodge filtration of the twisted connection

$$\nabla := d + d(f^e)$$

on $U := X \setminus S$. Thus $f : U \to \mathbb{A}$ is proper. As before we let

$$\breve{\Omega}^i_X = \Omega^i_X(\log S).$$

In this case, the associated Hodge filtration $F^\lambda(\nabla)$ on the de Rham complex reads

$$\begin{tikzcd}
F^0(\nabla) & F^0(\Omega^\bullet_X(\log \nabla)) \arrow[l, "\sim"'] & \mathcal{O}_X(-S) \arrow[r] & \breve{\Omega}^1_X((e-1)S) \arrow[r] & \breve{\Omega}^2_X((2e-1)S) \\
F^{m/e}(\nabla) & & & \breve{\Omega}^1_X((e-m)S) \arrow[r] & \breve{\Omega}^2_X((2e-m)S) \\
F^{1+m/e}(\nabla) & & & & \breve{\Omega}^2_X((e-m)S)
\end{tikzcd}$$

where $m \in \{1, 2, \cdots, e\}$. The attached spectral sequence reads

$$E_1^{pq} = \mathbb{H}^{p+q}\left(X, \mathrm{Gr}^{p/e}(\nabla)\right) \Longrightarrow H^{p+q}_{\mathrm{dR}}(U, \nabla). \tag{22}$$

Lemma 3.1 *Let ω_S be the dualizing sheaf of S. Then we have natural isomorphisms*

$$\mathrm{Gr}^{1+m/e}(\nabla) \xrightarrow{\sim} \omega_S[-2] \qquad \text{for } m = 1, 2, \cdots, e-1. \tag{23}$$

Proof. For any integer n, we have the adjunction exact sequence

$$0 \to \breve{\Omega}^2((n-1)S) \to \breve{\Omega}^2(nS) \to \omega_S(nS) \to 0$$

and the isomorphism

$$\omega_S(nS) \xrightarrow{\text{mult. by } f^{-n}} \omega_S$$

(between the two sheaves on X supported on S). The assertion then follows. $\square$

Lemma 3.2 *We have*

$$H^1(X, \mathcal{O}(-S)) = 0 = H^0(X, \breve{\Omega}^1).$$

Proof. We have the short exact sequence of sheaves on X

$$0 \to \mathcal{O}(-S) \to \mathcal{O} \to \mathcal{O}_S \to 0$$

which yields the exact

$$0 = H^0(X, \mathcal{O}(-S)) \to H^0(X, \mathcal{O}) \to H^0(S, \mathcal{O}_S) \to H^1(X, \mathcal{O}(-S)) \to H^1(X, \mathcal{O}).$$

Since S is connected, the complex vector space $H^0(S, \mathcal{O}_S)$ is one-dimensional. Since $H^1(X, \mathcal{O}) = 0$ by assumption, we have $H^1(X, \mathcal{O}(-S)) = 0$.

Let $\widetilde{S}$ be the normalization of S. Consider the exact

$$0 \to \Omega^1 \to \breve{\Omega}^1 \xrightarrow{\text{res.}} \mathcal{O}_{\widetilde{S}} \to 0$$

which gives

$$0 = H^0(X, \Omega^1) \to H^0(X, \breve{\Omega}^1) \to H^0(\widetilde{S}, \mathcal{O}_{\widetilde{S}}) \xrightarrow{c} H^1(X, \Omega^1).$$

The negative-definiteness of the intersection pairing on the irreducible components of S implies that the cycle map c is injective. Hence we have $H^0(X, \breve{\Omega}^1) = 0$. □

Proposition 3.3 *The short exact sequences*

$$0 \to F^{(m+1)/e}(\nabla) \to F^{m/e}(\nabla) \to \mathrm{Gr}^{m/e}(\nabla) \to 0$$

induce short exact sequences

$$0 \to \mathbb{H}^1\left(X, F^{(m+1)/e}(\nabla)\right) \to \mathbb{H}^1\left(X, F^{m/e}(\nabla)\right) \to \mathbb{H}^1\left(X, \mathrm{Gr}^{m/e}(\nabla)\right) \to 0$$

for $m = 0, 1, \cdots, e$, i.e. in the attached spectral sequence, $E_1^{p,1-p} = E_\infty^{p,1-p}$.

Proof. For $m = 0$, the assertion follows by the lemma above and the fact that

$$H^0\left(X, \mathrm{Gr}^0(\nabla)\right) = H^0(X, \mathcal{O}(-S)) = 0.$$

For $m = e$, we have

$$0 = \mathbb{H}^1\left(X, F^{1+}(\nabla)\right) \to \mathbb{H}^1\left(X, F^1(\nabla)\right) \to \mathbb{H}^1\left(X, \mathrm{Gr}^1(\nabla)\right)$$

with $\mathrm{Gr}^1(\nabla) \cong [0 \to \breve{\Omega}^1 \to \omega_S]$. Since $\mathbb{H}^1\left(X, \mathrm{Gr}^1(\nabla)\right)$ is a subset of $H^0(X, \breve{\Omega}^1)$, the assertion follows again by the lemma above.

Now assume that $1 < m < e$. We have

$$\begin{aligned} 0 = \mathbb{H}^1\left(X, \mathrm{Gr}^{m/e}(\nabla)\right) &\to \mathbb{H}^1\left(X, F^{(m+1)/e}(\nabla)\right) \\ &\to \mathbb{H}^1\left(X, F^{m/e}(\nabla)\right) \xrightarrow{\alpha} \mathbb{H}^1\left(X, \mathrm{Gr}^{m/e}(\nabla)\right). \end{aligned} \tag{24}$$

We need to show that the map α is surjective. By duality and (23), the dimension $h^1\left(X, \mathrm{Gr}^{m/e}(\nabla)\right)$ of the last term equals

$$h^3\left(X, \mathrm{Gr}^{2-m/e}(\nabla^\vee)\right) = h^1(S, \omega_S) = 1.$$

On the other hand, it is clear that the non-zero differential form $d(f^{e-m})$, with pole order exactly $(e - m + 1)$ along S, lies in

$$\mathbb{H}^1\left(X, F^{m/e}(\nabla)\right) = \ker\left\{\Gamma\left(X, \breve{\Omega}^1((e-m)S)\right) \xrightarrow{\nabla = d + ef^{e-1}df} \Gamma\left(X, \breve{\Omega}^2((2e-m)S)\right)\right\}$$

but is not in the subspace $\mathbb{H}^1\left(X, F^{(m+1)/e}(\nabla)\right)$. Thus α in (24) must be surjective. □

Corollary 3.4 *The Hodge to de Rham spectral sequence* (22) *degenerates at E_1-terms in this case.*

Proof. First we have $H^0_{\mathrm{dR}}(X,\nabla)\subset H^0(X,\mathcal{O}(-S))=0$. Together with the proposition above, we have

$$h^i_{\mathrm{dR}}(U,\nabla)=\sum_{0\le\lambda\le 2}h^i\left(U,\mathrm{Gr}^\lambda(\nabla)\right) \tag{25}$$

for $i=0,1$. By duality the equality (25) also holds for $i=3,4$, which then forces (25) to hold for $i=2$. This completes the proof. $\square$

Remark. In this case, we have the Hodge symmetry

$$\dim H^i\left(U,\mathrm{Gr}^\lambda(\nabla)\right)=\dim H^i\left(U,\mathrm{Gr}^{2-\lambda}(\nabla)\right).$$

4 The hypersurface case

We first recall the following well-known relation between the exponential sums and counting solutions of equations over a finite field. Let f be a regular function on a quasi-projective variety U over a finite field κ. To count the number $N(f)$ of the zero set

$$\{x\in U(\kappa)\mid f(x)=0\},$$

one brings in a non-trivial additive character $\chi:\kappa\to\mathbb{C}^\times$ and introduces a new variable $z=$ a fixed coordinate of an affine line $\mathbb{A}$ over κ. Let $\widetilde{f}=zf$, a regular function on $\mathbb{A}\times U$. Then we have

$$\sum_{x\in(\mathbb{A}\times U)(\kappa)}\chi\left(\widetilde{f}(x)\right)=q\cdot N(f)$$

where q is the cardinality of $\kappa=\mathbb{A}(\kappa)$.

Now the exponential sum in the left hand term of the equality above is related to the finite-field counterpart of the twisted de Rham cohomology while the right hand term consists of information of the closed subscheme of U defined by f. This suggests that in the world over the field of complex numbers, the de Rham cohomology of the connection $\nabla_{\widetilde{f}}$ over the product $\mathbb{A}\times U$ together with its irregular Hodge filtration should reflect the usual de Rham cohomology of the closed subscheme (f) defined by f with the usual Hodge filtration. We work out this analogue in this section. We consider the case where $(f)_{\mathrm{red}}$ defines a smooth divisor of U since we only have defined the twisted de Rham complexes and the filtrations for smooth varieties.

Lemma 4.1 *Let U be quasi-projective and smooth and $\widetilde{U}=\mathbb{A}\times U$. Consider the two projections*

$$\mathbb{A}\xleftarrow{a}\widetilde{U}\xrightarrow{b}U.$$

(i) *Let ∇ be the twisted connection on $\mathbb{A}$ associated with the identity map. Let $\widetilde{\nabla}=a^*\nabla=\nabla\boxtimes d$ be the pullback connection on $\widetilde{U}$. Then for any i,λ we have*

$$H^i\left(\widetilde{U},F^\lambda(\widetilde{\nabla})\right)=0.$$

(ii) *Let ∇ be the twisted connection associated with a regular function on U and $\widetilde{\nabla} = b^*\nabla = d \boxtimes \nabla$ be the pullback. Then for any i, λ we have*

$$H^i\left(\widetilde{U}, F^\lambda(\widetilde{\nabla})\right) = H^i\left(U, F^\lambda(\nabla)\right).$$

Proof. Let (X, S) be a good compactification of (U, f) where $f = 0$ in (i) or the regular function in (ii). Then $(\mathbb{P} \times X, \{\infty\} \times X \cup \mathbb{P} \times S)$ is in fact a good compactification of $(\widetilde{U}, \mathrm{id} \circ a)$ in (i) or $(\widetilde{U}, f \circ b)$ in (ii).

(i) Let $F^\lambda_\boxtimes$ be the exterior product filtration on $\mathbb{P} \times X$ of $F^i(\nabla)$ on $\mathbb{P}$ and $F^j(d)$ on X. By Cor.1.2 the natural inclusion $F^\lambda_\boxtimes \to F^\lambda(\widetilde{\nabla})$ is a quasi-isomorphism. Thus we only need to compute the hypercohomology of $F^\lambda_\boxtimes$. On the other hand, on the good compactification $\mathbb{P}$ of $\mathbb{A}$ we have

$$\mathrm{Gr}^0(\nabla) \cong \mathcal{O}_\mathbb{P}(-1) \quad \text{and} \quad F^{0+}(\nabla) = \breve{\Omega}^1_{\mathbb{A}\subset\mathbb{P}}[-1] \cong \mathcal{O}_\mathbb{P}(-1)[-1].$$

Thus $F^\lambda_\boxtimes$ is quasi-isomorphic to an extension of

$$A := \mathrm{Gr}^0(\nabla) \boxtimes F^\lambda(d) \cong \mathcal{O}_\mathbb{P}(-1) \boxtimes F^\lambda(d)$$

by

$$B := F^{0+}(\nabla) \boxtimes F^{\lambda-}(d) \cong \mathcal{O}_\mathbb{P}(-1)[-1] \boxtimes F^\lambda(d).$$

Since A and B have trivial hypercohomology, the assertion follows.

(ii) Similarly let $F^\lambda_\boxtimes$ be the product filtration on $\mathbb{P} \times X$ of $F^i(d)$ on $\mathbb{P}$ and $F^j(\nabla)$ on X. We have the quasi-isomorphism $F^\lambda_\boxtimes \xrightarrow{\sim} F^\lambda(\widetilde{\nabla})$ by Prop.1.1. This time on $\mathbb{P}$ we have

$$\mathrm{Gr}^0(\nabla) \cong \mathcal{O}_\mathbb{P} \quad \text{and} \quad F^{0+}(\nabla) = \breve{\Omega}^1_{\mathbb{A}\subset\mathbb{P}}[-1] \cong \mathcal{O}_\mathbb{P}(-1)[-1].$$

Thus $F^\lambda_\boxtimes$ on $\mathbb{P} \times X$ has the same hypercohomology as $F^\lambda(\nabla)$ on X. □

Lemma 4.2 *Let f be a nowhere vanishing regular function on a smooth quasi-projective U° and $\widetilde{f} = zf$ on $\mathbb{A} \times U^\circ$ where $z =$ identity on $\mathbb{A}$. Let $\nabla = d + d\widetilde{f}$ be the twisted connection on $\mathbb{A} \times U^\circ$. Then for all i, λ we have*

$$H^i\left(\mathbb{A} \times U^\circ, F^\lambda(\nabla)\right) = 0.$$

Proof. We have the commutative diagram

$$\begin{tikzcd}
\mathbb{A} \times U^\circ \arrow[r, "\alpha"] \arrow[dr, "\widetilde{f}=zf"'] & \mathbb{A} \times U^\circ \arrow[d, "\mathrm{pr}_1"] \\
 & \mathbb{A}
\end{tikzcd}$$

where

$$\alpha(z, x) := (zf(x), x),$$

is an isomorphism. Thus to prove the assertion, one reduces to the case where $\widetilde{f} = z$ via the isomorphism α. The assertion then follows from Lemma 4.1(i). □

Theorem 4.3 *Consider a pair* (U,f) *as before. Let* $V=(f)_{\mathrm{red}}$ *be the closed subvariety of* U *defined by* f. *Let* $\widetilde{f}=zf$ *on* $\mathbb{A}\times U$ *where* $z=$ *identity on* $\mathbb{A}$. *Assume that* V *is smooth. Then, for any* i,λ,

$$H^{i+2}_{\mathrm{dR},*}\left(\mathbb{A}\times U,\nabla_{\widetilde{f}}\right)=H^{i}_{\mathrm{dR},*}(V)$$

and

$$H^{i+2}_{*}\left(\mathbb{A}\times U,\mathrm{Gr}^{\lambda+1}(\nabla_{\widetilde{f}})\right)=H^{i-\lceil\lambda\rceil}_{*}\left(V,\Omega^{\lceil\lambda\rceil}\right)$$

where $*=\mathrm{c}$ *or nothing.*

Proof. By duality, it is enough to consider the case for the cohomology with compact support.

Let $U^\circ=U\setminus V$. Thus the three

$$\mathbb{A}\times U^\circ\hookrightarrow\mathbb{A}\times U\hookleftarrow\mathbb{A}\times V$$

form an open-closed decomposition and, by Prop.1.10, we have the long exact sequence

$$\cdots\to H^i_{\mathrm{c}}\left(\mathbb{A}\times U^\circ,F^\lambda(\nabla)\right)\to H^i_{\mathrm{c}}\left(\mathbb{A}\times U,F^\lambda(\nabla)\right)\to H^i_{\mathrm{c}}\left(\mathbb{A}\times V,F^\lambda(d)\right)\to\cdots.$$

By the dual of the lemma above and the Künneth formula, we then have

$$H^i_{\mathrm{c}}\left(\mathbb{A}\times U,F^\lambda(\nabla)\right)=H^i_{\mathrm{c}}\left(\mathbb{A}\times V,F^\lambda(d)\right)=H^{i-2}_{\mathrm{c}}\left(V,F^{\lceil\lambda\rceil-1}(d)\right).$$

The assertions now follow. □

Remark. The above theorem implies in particular that the Hodge to de Rham spectral sequence degenerates at E_1-terms in the case $\nabla=\nabla_{\widetilde{f}}$. The fact that the filtration $F^\lambda_{\mathrm{c}}(d)$ indeed induces the Hodge filtration of the canonical mixed Hodge structure on $H_{\mathrm{dR,c}}(V)$ can be found in [10, §4.3.3 and Prop.4.3.6].

Using the same idea, one has the following statements, also motivated by counting the number of the solutions of equations over finite fields.

Corollary 4.4 *Let* $f_1,\cdots,f_n$ *be regular functions on* U *and consider* $\widetilde{f}=\sum_{i=1}^n z_if_i$ *on* $\mathbb{A}^n\times U$ *where* $\{z_i\}=$ *the cartesian coordinates on* $\mathbb{A}^n$. *Suppose that each* $(f_i)_{\mathrm{red}}$ *is smooth and* $\sum_{i=1}^n(f_i)_{\mathrm{red}}$ *is a strict normal crossing divisor. Let* $W=\bigcap_{i=1}^n(f_i)_{\mathrm{red}}$. *Then, for any* j,λ,

$$H^{j+2n}_{\mathrm{dR},*}\left(\mathbb{A}^n\times U,\nabla_{\widetilde{f}}\right)=H^{j}_{\mathrm{dR},*}(W)$$

and

$$H^{j+2n}_{*}\left(\mathbb{A}^n\times U,\mathrm{Gr}^{\lambda+n}(\nabla_{\widetilde{f}})\right)=H^{j-\lceil\lambda\rceil}_{*}\left(W,\Omega^{\lceil\lambda\rceil}\right)$$

where $*=\mathrm{c}$ *or nothing.*

Proof. Again by duality, it is enough to consider $* = \mathrm{c}$. The above theorem gives the results for $n = 1$.

In general, one considers the commutative diagram

$$\begin{tikzcd}
\mathbb{A}^n \times V \arrow[r] & \mathbb{A}^n \times U & \mathbb{A}^n \times U^\circ \arrow[l] \arrow[d, "\alpha"'] \arrow[dr, "\widetilde{f}|_{\mathbb{A}^n \times U^\circ}"] & \\
 & & \mathbb{A}^n \times U^\circ \arrow[r, "\mathrm{pr}_1"'] & \mathbb{A}
\end{tikzcd}$$

where $V = (f_1)_{\mathrm{red}}$, $U^\circ = U \setminus V$ and

$$\alpha(z_1, \cdots, z_n; x) := (z_1 f_1(x) + \cdots + z_n f_n(x), z_2, \cdots, z_n; x)$$

defines an isomorphism. Now the triangle in the diagram shows that on $\mathbb{A}^n \times U^\circ$, the connection is isomorphic to $d + dz_1$. Then by Lemma 4.2 (applied to $f = 1$ on $\mathbb{A}^{n-1} \times U^\circ$), the cohomology of this connection and of its filtered pieces all vanish. Therefore the long exact sequence associated with the open-closed decomposition in the upper row of the diagram gives

$$H^{j+2}_{\mathrm{dR,c}}\left(\mathbb{A}^n \times U, F^\lambda(\nabla)\right) = H^{j+2}_{\mathrm{dR,c}}\left(\mathbb{A}^n \times V, F^\lambda(\nabla|_{\mathbb{A}^n \times V})\right) = H^{j}_{\mathrm{dR,c}}\left(\mathbb{A}^{n-1} \times V, F^{\lambda-1}(\nabla')\right)$$

for any j, λ with $\nabla' = d + d(\sum_{i=2}^n z_i f_i)$ on $\mathbb{A}^{n-1} \times V$. Here the second equality follows from the fact that $\nabla|_{\mathbb{A}^n \times V} = d_{z_1} \boxtimes \nabla'$ and by the dual of Lemma 4.1(ii). The statements now follow by induction on the number n of the defining equations of W. $\square$

5 The toric case

Suppose U is a torus. Inspired by the investigation [1] of exponential sums over a torus via Dwork's p-adic methods and the work of Kouchnirenko [20] on the Milnor numbers of isolated singularities, Adolphson and Sperber in [2] study the twisted de Rham cohomology on U (in fact in a more general setting which also allows multiplicative twists). They derive that for generic f, the twisted de Rham cohomology is concentrated in a single degree. The method there is to introduce a filtration, already appeared in [20], on the de Rham chain complex and show that the associated graded complex has non-trivial cohomology only at one degree. In this section we recall their filtration and show that the induced filtration on the de Rham cohomology coincides with our irregular Hodge filtration for f generic.

Our reference for the theory of toric variety is [11]. In particular, see [11, p.48] for the existence of the equivariant resolution of singularities and [11, p.61] for the computation of the valuation of a function on a toric divisor.

In this section we let

$$U = (\mathbb{A} \setminus 0)^n \quad \text{and} \quad f \in \mathcal{O}(U) = \mathbb{C}[x^{\pm 1}]$$

where $x = (x_1, \cdots, x_n)$ is the system of cartesian coordinates of U. Again let $\nabla = d + df$ be the twisted connection on U. Recall the following.

Definition. Write $f = \sum_{\alpha \in \mathbb{Z}^n} c(\alpha) x^\alpha$.

(i) The *Newton polyhedron* $\Delta(f)$ of f is the convex hull in $\mathbb{R}^n$ of the finite set

$$\{0\} \cup \{\alpha \in \mathbb{Z}^n \,|\, c(\alpha) \neq 0\}.$$

(ii) The function f is called *non-degenerate with respect to* $\Delta(f)$ if for any face δ of $\Delta(f)$ with $0 \notin \delta$, the system of equations

$$f_\delta = \frac{\partial f_\delta}{\partial x_1} = \cdots = \frac{\partial f_\delta}{\partial x_n} = 0 \tag{26}$$

has no solution on U where $f_\delta := \sum_{\alpha \in \delta} c(\alpha) x^\alpha$.

One regards $\Delta(f)$ as sitting in the space of characters $M_\mathbb{R} := \mathbb{R} \underset{\mathbb{Z}}{\otimes} \operatorname{Hom}(U, \mathbb{C}^\times)$. It then defines a fan on the dual space $N_\mathbb{R} := \operatorname{Hom}_\mathbb{R}(M, \mathbb{R})$ where each codimension one face of $\Delta(f)$ corresponds to a ray in the fan, pointing to the inward normal direction with respect to the natural pairing $N_\mathbb{R} \times M_\mathbb{R} \to \mathbb{R}$. Now one can refine and enlarge the fan to make a cone decomposition of $N_\mathbb{R}$ such that the associated toric variety X_{tor} is smooth and proper and the toric boundary $S := X_{\mathrm{tor}} \setminus U$ is a simple normal crossing divisor of X_{tor}. Each ray in this refined fan corresponds to an irreducible component of S. We fix this X_{tor} in the sequel. We have the commutative diagram

$$\begin{tikzcd}
U \arrow[r, "f"] \arrow[d] & \mathbb{A} \arrow[d] \\
X_{\mathrm{tor}} \arrow[r, dashed] & \mathbb{P}
\end{tikzcd}$$

where the two vertical arrows are the inclusions but the lower arrow is just a rational function on X_{tor} in general.

The connection ∇ on U again extends to the complex on X_{tor}

$$(\Omega^\bullet(*S), \nabla) = \Big[\mathcal{O}(*S) \xrightarrow{\nabla} \Omega^1(*S) \to \cdots \to \Omega^n(*S)\Big]$$

and we have

$$H^i_{\mathrm{dR}}(U, \nabla) = \mathbb{H}^i\big(X_{\mathrm{tor}}, (\Omega^\bullet(*S), \nabla)\big).$$

If $\dim \Delta(f) < n$, there is a decomposition $U = U' \times U''$ of U into two tori and $f' \in \mathcal{O}(U')$ such that $f = f' \circ \mathrm{pr}_{U'}$. In this case $\nabla = \nabla_{f'} \boxtimes d$ and our discussion of the irregular Hodge filtration also reduces to the product situation. Thus for simplicity, we will assume that $\dim \Delta(f) = n$ in the rest of this section. The general case then can be deduced easily.

(a) *The Newton polyhedron filtration*

We define the *Newton polyhegron filtration* $F^{\lambda}_{\mathrm{NP}}(\nabla)$ of $(\Omega^{\bullet}(*S),\nabla)$ on X_{tor} similar to the filtration $F^{\lambda}(\nabla)$ for a good compactification X. Again let P be the pole divisor of f on X_{tor}. Let

$$F^{\lambda}_{\mathrm{NP}}(\nabla) := \Big[\mathcal{O}(\lfloor -\lambda P\rfloor) \xrightarrow{\nabla} \breve{\Omega}^1(\lfloor (1-\lambda)P\rfloor) \to \cdots \to \breve{\Omega}^p(\lfloor (p-\lambda)P\rfloor) \to \cdots\Big]^{\geq \lceil \lambda \rceil}. \tag{27}$$

Notice that if the origin is contained in the interior of $\Delta(f)$, then the morphism $f : U \to \mathbb{A}$ is proper and the filtration $F^{\lambda}_{\mathrm{NP}}(\nabla)$ is indeed exhaustive.

To compute the hypercohomology of $F^{\lambda}_{\mathrm{NP}}(\nabla)$, first notice that on the toric variety X_{tor} the locally free sheaf $\breve{\Omega}^p$ is trivial for any p. Indeed as an $\mathcal{O}$-module it is globally generated by

$$\bigwedge^{p}\left\{\frac{dx_1}{x_1},\cdots,\frac{dx_n}{x_n}\right\}.$$

On the other hand for $p \geq \lambda$, we have [11, Prop.p.68 and Cor.p.74]

$$H^i\big(X_{\mathrm{tor}},\mathcal{O}(\lfloor (p-\lambda)P\rfloor\big) = \begin{cases} \begin{array}{l}\mathbb{C}\text{-vector space generated by } \{x^{\alpha}\}\\ \text{where } \alpha \text{ runs over the lattice points}\\ \text{inside the dilated polyhedron } (p-\lambda)\cdot\Delta(f)\end{array} & \text{if } i=0 \\ 0 & \text{if } i\neq 0. \end{cases}$$

Thus one obtains

$$\mathbb{H}^i\left(X_{\mathrm{tor}}, F^{\lambda}_{\mathrm{NP}}(\nabla)\right) = H^i\left(\Gamma\left(X_{\mathrm{tor}}, F^{\lambda}_{\mathrm{NP}}(\nabla)\right)\right) \tag{28}$$

and this cohomology does not depend on the choice of X_{tor}.

Theorem 5.1 *Suppose* $\Delta(f) = n$ *and* f *is non-degenerate with respect to* $\Delta(f)$. *With notations as above, we have*

(i) *For* $\lambda \leq 0$ *the inclusion* $F^{\lambda}_{\mathrm{NP}}(\nabla) \to (\Omega^{\bullet}(*S),\nabla)$ *on* X *is a quasi-isomorphism.*

(ii) $H^i\left(\Gamma\left(X_{\mathrm{tor}}, \mathrm{Gr}^{\lambda}_{\mathrm{NP}}(\nabla)\right)\right) \neq 0$ *only if* $i = n$.

(iii) *Let* $\mathrm{Vol}(f)$ *be the usual Euclidean volume of* $\Delta(f)$ *in* $\mathbb{R}^n$. *Then*

$$\dim H^i_{\mathrm{dR}}(U,\nabla) = \begin{cases} n!\cdot\mathrm{Vol}(f) & \textit{if } i=n \\ 0 & \textit{otherwise.}\end{cases}$$

The proof of (i) is established in [2, pp.70-73] where the authors show the quasi-isomorphism between the two complexes $(\hat{K}^{\cdot},\delta_{f,\alpha})$ and $(K^{\cdot}_0,\delta_{f,\alpha})$, which correspond to $F^{0}_{\mathrm{NP}}(\nabla)$ and $(\Omega^{\bullet}(*S),\nabla)$, respectively. For (ii), see [2, (4.3)], cf. [1, Thm.2.14] and [20, Th.2.8]. For (iii), see [2, Thm1.4 and Thm.4.1].[1]

Combined with (28), the above theorem implies the following.

---

[1] The relations between the notations here and in [2] are that

$$\Gamma\left(X_{\mathrm{tor}}, F^{\lambda}_{\mathrm{NP}}(\nabla)^{l}\right) = F_{-\lambda}\hat{K}^{l} \quad \text{and} \quad \mathrm{Gr}^{\bullet}_{\mathrm{NP}}(\nabla) = (\bar{K}^{\cdot},\bar{\delta}_{f,\alpha}).$$

Notice that there is a typo in [2, p.68]. In line 18, the weight $(k/e)-l$ should be $(k/e)+l$.

Corollary 5.2 *Suppose* $\Delta(f) = n$ *and* f *is non-degenerate with respect to* $\Delta(f)$*. The spectral sequence attached to the filtration* $F^{\lambda}_{\mathrm{NP}}(\nabla)$ *on* X_{tor} *converges to* $H_{\mathrm{dR}}(U, \nabla)$ *and degenerates at the initial stage.*

(*b*) *The comparison*

As already mentioned, the rational function f on X_{tor} is not yet a morphism to $\mathbb{P}$ in general and hence (X_{tor}, S) is not a good compactification of (U, f) for defining the irregular Hodge filtration. This is because the zero divisor Z and the pole divisor P of f intersect and one needs to perform blowups, say $\pi : X \to X_{\mathrm{tor}}$, in order to eliminate the indeterminacy and to obtain a morphism $f : X \to \mathbb{P}$. However when f is non-degenerate with respect to $\Delta(f)$, we can say more.

Proposition 5.3 *Suppose that* f *is non-degenerate with respect to* $\Delta(f)$*. Then on* X_{tor} *the zero divisor* Z *and the support of the pole divisor* $(P)_{\mathrm{red}}$ *of* f *intersect transversally and the intersections of* Z *with various toric strata of* $(P)_{\mathrm{red}}$ *are smooth.*

Proof. A codimension r toric stratum D of $(P)_{\mathrm{red}}$ is a dense torus sitting in an irreducible component of the intersection of certain irreducible components $D_1, \cdots, D_r$ of S. Each D_i corresponds to a ray in $N_{\mathbb{R}}$, which then corresponds to a face δ_i of $\Delta(f)$ (containing the exponents $\alpha \in \Delta(f)$ with most negative product with the direction of the ray). A face δ in the intersection of δ_i then corresponds to D and f_δ is the most singular term of the function f restricted to D since those monomials in f_δ are among the terms in f which have the highest pole order along D. We have $0 \notin \delta$ since otherwise f has no pole along D. Also the indeterminacy locus $Z \cap D$ on D is exactly the zero set defined by $f_\delta = 0$. Now the condition of emptiness of the solution of (26) (which becomes the usual Jocabian criterion after a change of variables) exactly says that $Z \cap D$ is smooth, which is what we want. $\square$

From now on we assume that f is non-degenerate with respect to $\Delta(f)$.

We construct one particular $\pi : X \to X_{\mathrm{tor}}$ to obtain $f : X \to \mathbb{P}$ as follows. One picks an irreducible component D of $Z \cap (P)_{\mathrm{red}}$ and then take the blowup X' along D. If the exceptional divisor E contributes to the pole of f on X', we perform the blowup along $E \cap Z'$ where Z' is the zero divisor of f on X'. Continue this procedure until f extends to a morphism to $\mathbb{P}$ along the exceptional locus on $X^{(k)}$. Let $Z^{(k)}$ and $P^{(k)}$ be the zero and pole divisors of f on $X^{(k)}$. Then one picks one irreducible of $Z^{(k)} \cap (P^{(k)})_{\mathrm{red}}$ and performs a sequence of blowups again as above. Repeating the procedure, one then obtains the commutative diagram

$$\begin{tikzcd}
X \arrow[r] \arrow[rrrrr, bend left, "\pi"] \arrow[rrrrrd, bend right, "f"'] & \cdots \arrow[r] & X_2 \arrow[r, "\varepsilon"] & X_1 \arrow[r] & \cdots \arrow[r] & X_{\mathrm{tor}} \arrow[d, dashed, "f"] \\
 & & & & & \mathbb{P}
\end{tikzcd} \tag{29}$$

where each step is the blowup along a smooth irreducible of the intersection of the zero and pole divisors of f.

Now $(X, X \setminus U)$ defines a good compactification of (U, f) and we have the filtration $F^{\lambda}(\nabla)$ on X. We shall show that there is a natural quasi-isomorphism $\mathcal{R}\pi_* F^{\lambda}(\nabla) \to F^{\lambda}_{\mathrm{NP}}(\nabla)$ on X_{tor} for each λ. Consequently they define the same filtration on $H_{\mathrm{dR}}(U, \nabla)$ and furthermore the Hodge to de Rham spectral sequence degenerates in this case.

For this and to simplify the notations, we consider the filtrations, called $F^{\lambda}_1(\nabla)$ and $F^{\lambda}_2(\nabla)$, of the twisted de Rham complexes on X_1 and X_2, respectively where $\varepsilon : X_2 \to X_1$ appears in the above sequence of blowups. They are defined exactly as in (27) (which does not require the variety is toric). Now notice that for $X_2 = X$ the filtration $F^{\lambda}_2(\nabla)$ is $F^{\lambda}(\nabla)$ for the good compactification X while for $X_1 = X_{\mathrm{tor}}$ the filtration $F^{\lambda}_1(\nabla)$ is the Newton polyhedron filtration $F^{\lambda}_{\mathrm{NP}}(\nabla)$ on the toric X_{tor}. One then checks immediately that

$$F^{\lambda}_2(\nabla) \subset \varepsilon^* \left(F^{\lambda}_1(\nabla) \right). \tag{30}$$

We want to understand the difference of this inclusion.

We look at the local situation over a point of the center of blowup in X_1. Prop.5.3 ensures the following. We can take $X_1 = \mathbb{D}^n$ with coordinates

$$\{x, y_1, \cdots, y_k, t_1, \cdots, t_l, z, \tau_1, \cdots, \tau_m\}$$

and $U = (\mathbb{D}^{\circ})^{1+k+l} \times \mathbb{D}^{1+m}$ with the boundary $S_1 = (xyt)$. The regular function on U is

$$f = \frac{z}{x^e y^r} = \frac{z}{x^e y_1^{r_1} \cdots y_k^{r_k}} \quad \text{(for some } e, r > 0\text{)}.$$

The center Ξ of blowup ε is given by $x = 0 = z$.

The blowup $X_2 \subset X_1 \times \mathbb{P}$ is given by the equation

$$xu = zv \quad ([u, v] \in \mathbb{P}).$$

We use the notations in the illustration of $X_2 \xrightarrow{\varepsilon} X_1$ below.



Here the exceptional divisor E is a split $\mathbb{P}^1$-bundle over the y-t-τ-coordinate plane Ξ of X_1. Let

$$e' := e - 1.$$

Thus the pole divisors of f on X_1 and X_2 are given by

$$P_1 = eA + rB \quad \text{and} \quad P_2 = eA + e'E + rB,$$

respectively. We have the information at the two points a_1 and a_2 in the table below with $\bar{v} = \frac{v}{u}$ and $\bar{u} = \frac{u}{v}$.

	a_1	a_2
coordinates	$\{\bar{v}, z, y, t, \tau\}$	$\{\bar{u}, x, y, t, \tau\}$
f	$\frac{1}{\bar{v}^e z^{e'} y^r}$	$\frac{\bar{u}}{x^{e'} y^r}$

We consider the refinement of the inclusion (30) as follows. Write $p = \lceil \lambda \rceil$. For $q = 0, 1, \cdots, (n-p)$, let $R_\lambda(q) = (R_\lambda(q)^\bullet, \nabla)$ be the complex on X_2 given by

$$R_\lambda(q)^{p+j} = \begin{cases} 0 & \text{if } j \leq 0 \\ \breve{\Omega}^{p+j}\big((\mu + je)A + (\mu + je)E + (\nu + jr)B\big) & \text{if } 0 \leq j \leq q \\ \breve{\Omega}^{p+j}\big((\mu + je)A + (\mu + je' + q)E + (\nu + jr)B\big) & \text{if } j > q \end{cases} \tag{31}$$

where $\mu = \lfloor (p - \lambda)e \rfloor$ and $\nu = \lfloor (p - \lambda)r \rfloor$. We have

$$R_\lambda(q-1)^{\leq p+q-1} = R_\lambda(q)^{\leq p+q-1} \tag{32}$$

and

$$R_\lambda(-1) := F_2^\lambda(\nabla) \subset R_\lambda(0) \subset R_\lambda(1) \subset \cdots \subset R_\lambda(n-p) = \pi^*\left(F_1^\lambda(\nabla)\right). \tag{33}$$

Notice that $R_\lambda(-1) = R_\lambda(0)$ if $\lfloor (p-\lambda)e \rfloor = \lfloor (p - \lambda)e' \rfloor$.

To understand the successive quotients of (33), we introduce one more complex. Let $S_2 = X_2 \backslash U$. For three integers ρ, η, ξ, we let $(K^\bullet_{\rho,\eta,\xi}, \nabla)$ be the subcomplex of $(\Omega^\bullet(*S_2), \nabla)$ on X_2 whose degree-j term is given by

$$K^j_{\rho,\eta,\xi} = \breve{\Omega}^j\left((\rho + je)A + (\eta + je')E + (\xi + jr)B\right).$$

One has

$$R_\lambda(q)^{\geq p+q} = \left(K^\bullet_{\mu - pe, \mu - pe' + q, \nu - pr}, \nabla\right)^{\geq p+q}. \tag{34}$$

Lemma 5.4 *The inclusion*

$$\left(K^\bullet_{\rho,\eta,\xi}, \nabla\right) \subset \left(K^\bullet_{\rho,\eta+1,\xi}, \nabla\right)$$

is a quasi-isomorphism of complexes on X_2 for any $\rho, \eta, \xi \in \mathbb{Z}$ with the condition that $\eta \geq 0$ if $e' = 0$.

Proof. At the point a_1, one only needs to consider the case where $e = 1$, thanks to Prop.1.3. In this case, we have the exterior product decomposion

$$(K^\bullet_{\rho,\eta,\xi}, \nabla) = (K^\bullet_{\rho,\xi}, \nabla') \boxtimes \left[\frac{1}{z^\eta}\mathcal{O}_z \xrightarrow{d} \frac{1}{z^\eta}\breve{\Omega}^1_z\right]$$

where $K^\bullet_{\rho,\xi}$ is defined similarly as the definition of $K^\bullet_{\rho,\eta,\xi}$ above but now on the coordinates $\{\bar{v}, y, t, \tau\}$ for the connection ∇' attached to $1/(\bar{v}y^r)$. The assertion then follows from the fact that

$$\left[\mathcal{O}_z \xrightarrow{d} \breve{\Omega}^1_z\right] \to \left[\frac{1}{z^i}\mathcal{O}_z \xrightarrow{d} \frac{1}{z^i}\breve{\Omega}^1_z\right]$$

is a quasi-isomorphism for any $i \geq 0$.

The case for points between a_1 and a_2 is similar.

At the point a_2, let $\mathcal{O}$ be the coordinate ring and $\bar{\mathcal{O}} = \mathcal{O}/x\mathcal{O}$. One has to check the exactness of

$$\begin{aligned} &\frac{\breve{\Omega}^{j-1}((\eta+1-e')E+(\xi-r)B)}{\breve{\Omega}^{j-1}((\eta-e')E+(\xi-r)B)} \xrightarrow{\nabla_{j-1}} \frac{\breve{\Omega}^{j}((\eta+1)E+\xi B)}{\breve{\Omega}^{j}(\eta E+\xi B)} \\ &\qquad\qquad\qquad\qquad\qquad \xrightarrow{\nabla_{j}} \frac{\breve{\Omega}^{j+1}((\eta+1+e')E+(\xi+r)B)}{\breve{\Omega}^{j+1}((\eta+e')E+(\xi+r)B)} \end{aligned} \tag{35}$$

where now the connection is the $\bar{\mathcal{O}}$-linear map given by the left cup product with

$$\frac{1}{x^{e'}y^{r}}\left(d\bar{u} - \bar{u}\frac{e'dx}{x} - \bar{u}\sum_{i=1}^{k}\frac{r_i dy_i}{y_i}\right) = \frac{\bar{u}}{x^{e'}y^{r}}\left(\frac{d\bar{u}}{\bar{u}} - \frac{e'dx}{x} - \sum_{i=1}^{k}\frac{r_i dy_i}{y_i}\right).$$

First suppose that $e' \geq 1$. Let

$$\Lambda_i := \bar{\mathcal{O}} \cdot \bigwedge^{i}\left\{\frac{d\bar{u}}{u}, \frac{dx}{x}, \frac{dy}{y}, \frac{dt}{t}, d\tau\right\}$$

be the $\bar{\mathcal{O}}$-module generated by i-forms. Now notice that the complex

$$\frac{1}{\bar{u}^2 x^{\eta+1-2e'}y^{\xi-2r}}\Lambda_{j-2} \xrightarrow{\nabla_{j-2}} \frac{1}{\bar{u}x^{\eta+1-e'}y^{\xi-r}}\Lambda_{j-1} \xrightarrow{\nabla_{j-1}} \frac{1}{x^{\eta+1}y^{\xi}}\Lambda_j \xrightarrow{\nabla_j} \frac{\bar{u}}{x^{\eta+1+e'}y^{\xi+r}}\Lambda_{j+1},$$

being isomorphic to the Koszul complex associated with

$$\{v_\infty, v_0, v_i\}_{i=1}^{k} \quad \text{corresponding to } \frac{\bar{u}}{x^{e'}y^{r}}\left\{\frac{d\bar{u}}{\bar{u}}, \frac{-e'dx}{x}, \frac{-r_i dy_i}{y_i}\right\}_{i=1}^{k},$$

is exact. Thus if $\nabla_j(\alpha) = 0$ for some α in our degree-j piece, there exists a

$$\beta = d\bar{u}\wedge\beta_1 + \beta_2 \quad \text{with} \left\{\begin{array}{l} \beta_1 \in \frac{1}{\bar{u}^2 x^{\eta+1-e'}y^{\xi-r}}\bar{\mathcal{O}}\cdot\bigwedge^{j-2}\left\{\frac{dx}{x}, \frac{dy}{y}, \frac{dt}{t}, d\tau\right\} \\ \beta_2 \in \frac{1}{\bar{u}x^{\eta+1-e'}y^{\xi-r}}\bar{\mathcal{O}}\cdot\bigwedge^{j-1}\left\{\frac{dx}{x}, \frac{dy}{y}\frac{dt}{t}, d\tau\right\} \end{array}\right.$$

such that $\nabla_{j-1}(\beta) = \alpha$. By subtracting $\nabla_{j-2}(x^{e'}y^{r}\beta_1)$ to β, we may assume that $\beta_1 = 0$. Then the part $x^{-e'}y^{-r}d\bar{u}\wedge\beta_2$ of $\alpha = \nabla_{j-1}(\beta)$ does not have $\bar{u}$ in the denominator, and hence neither does β_2. Therefore $\beta \in \frac{1}{x^{\eta+1-e'}y^{\xi-r}}\Lambda_{j-1}$ and (35) is exact.

The case $e' = 0$ is similar. $\square$

Lemma 5.5 *Recall the complex $R_\lambda(q)$ on X_2 defined in* (31)*. For $q = 0, 1, \cdots, n-p$, the quotient $R_\lambda(q)/R_\lambda(q-1)$ is quasi-isomorphic to a direct sum of the relative degree (-1)-invertible sheaves $\mathcal{O}_{E/\Xi}(-1)$ concentrated at degree $p+q$.*

Proof. Let $\mu = \lfloor (p-\lambda)e \rfloor$ and $\nu = \lfloor (p-\lambda)r \rfloor$. Let $a = \mu - pe$, $b = \mu - pe' + q$ and $c = \nu - pr$. By the relations (32) and (34) and the previous lemma, we have

$$\frac{R_\lambda(q)}{R_\lambda(q-1)} \overset{\sim}{\longleftarrow} \ker \left\{ \frac{K^{p+q}_{a,b,c}}{K^{p+q}_{a,b-1,c}} \overset{\nabla}{\rightarrow} \frac{K^{p+q+1}_{a,b,c}}{K^{p+q+1}_{a,b-1,c}} \right\} \overset{\sim}{\longleftarrow} \mathrm{Image} \left\{ \frac{K^{p+q-1}_{a,b,c}}{K^{p+q-1}_{a,b-1,c}} \overset{\nabla}{\rightarrow} \frac{K^{p+q}_{a,b,c}}{K^{p+q}_{a,b-1,c}} \right\}. \tag{36}$$

(Notice that if $e' = 0$ and $b = 0$, then we have $q = \mu = 0$ and $R_\lambda(-1) = R_\lambda(0)$. So there is nothing to prove.)

Let $\eta = \mu + qe$ and $\xi = \nu + qr$. Away from $u = 0$ (neighborhood of a_1), the $\mathcal{O}_E$-module $K^{p+q-1}_{a,b,c}/K^{p+q-1}_{a,b-1,c}$ is generated by ω_1 of the four types listed in the table

$\bar{v}^{\eta-e} z^{\eta-e'} y^{\xi-r} \cdot \omega_1$	$x^{\eta-e'} y^{\xi-r} \cdot \omega_2$
$\zeta_1 \cdots \zeta_{p+q-1}$	$\zeta_1 \cdots \zeta_{p+q-1}$
$\frac{d\bar{v}}{\bar{v}} \zeta_1 \cdots \zeta_{p+q-2}$	$-\frac{d\bar{u}}{\bar{u}} \zeta_1 \cdots \zeta_{p+q-2}$
$\frac{dz}{z} \zeta_1 \cdots \zeta_{p+q-2}$	$\left(e\frac{dx}{x} + \sum_{i=1}^k r_i \frac{dy_i}{y_i} \right) \zeta_1 \cdots \zeta_{p+q-2}$
$\frac{d\bar{v}}{\bar{v}} \frac{dz}{z} \zeta_1 \cdots \zeta_{p+q-3}$	$\left(\frac{dx}{x} \sum_{i=1}^k r_i \frac{dy_i}{y_i} \right) \zeta_1 \cdots \zeta_{p+q-3}$

where

$$\{\zeta_j\} = \left\{ \frac{dy}{y}, \frac{dt}{t}, d\tau \right\}.$$

One checks that we have

$$\bar{u} \cdot \nabla(\omega_1) \equiv \nabla(\omega_2) \pmod{K^{p+q}_{a,b-1,c} = \breve{\Omega}^{p+q}\big(\eta A + (\eta - 1)E + \xi B\big)}$$

where ω_2 are the corresponding forms lying away from $v = 0$ (neighborhood of a_2) listed above. This equation shows that the last term of (36) is a direct sum of $\mathcal{O}_{E/\Xi}(-1)$. $\square$

Theorem 5.6 *Consider the pair (U, f) where U is a torus of dimension $n = \dim \Delta(f)$ and f is non-degenerate with respect to $\Delta(f)$. Then*

$$\dim H^i_{\mathrm{dR}}(U, \nabla) = \begin{cases} n! \cdot \mathrm{Vol}(f) & \text{if } i = n \\ 0 & \text{otherwise,} \end{cases}$$

the irregular Hodge filtration coincides with the filtration induced by $F^\lambda_{\mathrm{NP}}(\nabla)$ on any smooth toric compactification X_{tor} with simple normal crossing boundary $X_{\mathrm{tor}} \setminus U$, and the Hodge to de Rham spectral sequence degenerates at the initial stage.

Proof. We choose a good compactification X with $\pi : X \to X_{\mathrm{tor}}$ as constructed in (29). We have the natural inclusion

$$F^\lambda(\nabla) \subset \pi^* F^\lambda_{\mathrm{NP}}(\nabla) \tag{37}$$

for any $\lambda \in \mathbb{R}$ from (30). The refinement (33) and the above lemma show that there is a refinement of the inclusion (37) such that the successive quotients have trivial higher relative cohomology $\mathcal{R}^i \pi_*$ in any degree $i \geq 0$. Thus the adjoint

$$\mathcal{R}\pi_* F^\lambda(\nabla) \to F^\lambda_{\mathrm{NP}}(\nabla)$$

is indeed a quasi-isomorphism of complexes on X_{tor} for any λ. The assertions now follow from Thm.5.1 and Cor.5.2. $\square$

6 Relation with Sabbah's work

In this section, we first recall the work of Sabbah in [25] and apply the results to obtain the degeneracy of the Hodge to de Rham spectral sequence in the case where $f : X \to \mathbb{P}$ is a smooth morphism and $U = f^{-1}(\mathbb{A})$.

(a) *Results of Sabbah*

Fix a coordinate t of $\mathbb{A} \subset \mathbb{P}$ and let $z = t^{-1}$ be the local coordinate at $\infty \in \mathbb{P}$.

We start from a variation of pure Hodge structure $(\mathcal{M}, F, \theta)$ on $\mathbb{P}$. Here the triple $(\mathcal{M}, F, \theta)$ consists of

- a local free coherent $\mathcal{O}_{\mathbb{P}}$-module $\mathcal{M}$,
- the Hodge filtration F^λ, which is a finite filtration on $\mathcal{M}$ by locally free and co-free $\mathcal{O}_{\mathbb{P}}$-submodules with integral jumps satisfying

  $$\mathcal{M} = F^\lambda\mathcal{M} \oplus \overline{F^{w+1-\lambda}\mathcal{M}} \quad \text{for all } \lambda$$

  for a fixed integer w, and
- a connection $\theta : \mathcal{M} \to \Omega^1_{\mathbb{P}} \underset{\mathcal{O}_{\mathbb{P}}}{\otimes} \mathcal{M}$ satisfying the Griffiths transversality

  $$\theta\left(F^{\lambda+1}\mathcal{M}\right) \subset \Omega^1_{\mathbb{P}} \underset{\mathcal{O}_{\mathbb{P}}}{\otimes} F^\lambda\mathcal{M} \quad \text{for all } \lambda.$$

Let

$$\widetilde{\mathcal{M}} = \mathcal{M}(*\infty) := \mathcal{O}(*\infty) \underset{\mathcal{O}_{\mathbb{P}}}{\otimes} \mathcal{M}$$

and consider the meromorphic connection δ on $\widetilde{\mathcal{M}}$ defined by

$$\widetilde{\mathcal{M}} \xrightarrow{\delta := \theta + dt} \Omega^1_{\mathbb{P}} \underset{\mathcal{O}_{\mathbb{P}}}{\otimes} \widetilde{\mathcal{M}}. \tag{38}$$

In this setting, Sabbah in [25, §6] defines an exhaustive and separated filtration F^λ_{Del} on $\widetilde{\mathcal{M}}$ by locally free sheaves, which satisfies the Griffiths transversality with respect to δ. (This is called *Deligne's filtration* in the reference and thus the subscript in F^λ_{Del}.) The definition is as follows.

Away from ∞, the new filtration F^λ_{Del} coincides with the Hodge filtration F^λ on $\mathcal{M}$.

Let $\partial_z = \theta(d/dz) : \mathcal{M} \to \mathcal{M}$. Then at the point ∞, we define

$$F^\lambda_{\mathrm{Del}}\widetilde{\mathcal{M}} := \sum_{k \geq 0} \left(\partial_z - \frac{1}{z^2}\right)^k \frac{1}{z} F^{\lambda+k}\mathcal{M}. \tag{39}$$

(Notice that $\partial_z - 1/z^2 = \delta(d/dz)$.)

One checks that this yields an exhaustive and separated filtration of (38) by the two-term complexes

$$\left[F^\lambda_{\mathrm{Del}}\widetilde{\mathcal{M}} \xrightarrow{\delta} \Omega^1_{\mathbb{P}} \underset{\mathcal{O}_{\mathbb{P}}}{\otimes} F^{\lambda-1}_{\mathrm{Del}}\widetilde{\mathcal{M}}\right]. \tag{40}$$

Now by the Griffiths transversality of θ and induction on $k \geq 1$, an easy calculation shows that at ∞ we have

$$\left(\partial_z - \frac{1}{z^2}\right)^k \frac{1}{z} F^{\lambda+1}\mathcal{M} \equiv \frac{1}{z^{2k+1}} F^{\lambda+1}\mathcal{M} \quad (\mathrm{mod}\ \frac{1}{z^{2k-1}} F^{\lambda}\mathcal{M})$$

and hence one obtains

$$F^{\lambda}_{\mathrm{Del}}\widetilde{\mathcal{M}} = \sum_{k\geq 0} \frac{1}{z^{2k+1}} F^{\lambda+k}\mathcal{M}.$$

Globally one then has

$$F^{\lambda}_{\mathrm{Del}}\widetilde{\mathcal{M}} = \sum_{k\geq 0} \left(F^{\lambda+k}\mathcal{M}\right)((2k+1)\infty). \tag{41}$$

In particular if the jumps of the Hodge filtration F^{λ} are all non-negative integers, we have

$$F^{-\lambda}_{\mathrm{Del}}\widetilde{\mathcal{M}} = \left(F^{0}_{\mathrm{Del}}\widetilde{\mathcal{M}}\right)(2\lfloor\lambda\rfloor \cdot \infty) \quad \text{for any } \lambda \geq 0.$$

Notice that in this setting, the connection δ in (40) is an injection between Zariski sheaves on $\mathbb{P}$ since the $\mathcal{O}_{\mathbb{P}}$-linear factor dt in δ increases the pole order at ∞ by two while the $\mathbb{C}$-linear part θ increases the order by at most one.

Proposition 6.1 *Assume that the jumps of the Hodge filtration F^{λ} are all non-negative integers. Then for any $\lambda \leq 0$, the natural inclusion*

$$\left[F^{0}_{\mathrm{Del}}\widetilde{\mathcal{M}} \xrightarrow{\delta} \Omega^1_{\mathbb{P}} \otimes F^{-1}_{\mathrm{Del}}\widetilde{\mathcal{M}}\right] \to \left[F^{\lambda}_{\mathrm{Del}}\widetilde{\mathcal{M}} \xrightarrow{\delta} \Omega^1_{\mathbb{P}} \otimes F^{\lambda-1}_{\mathrm{Del}}\widetilde{\mathcal{M}}\right] \tag{42}$$

is a quasi-isomorphism of complexes on $\mathbb{P}$.

Proof. As before we refine the filtration F_{Del} such that the successive quotient complexes become easier to deal with. Let $\widetilde{F_{\mathrm{Del}}}^{\lambda}$ denote the complex in the right hand term of (42). For any positive integer i, consider the complex $\widetilde{F}^{-i}$ defined by

$$\widetilde{F}^{-i} = \left[\left(F^{0}_{\mathrm{Del}}\widetilde{\mathcal{M}}\right)(i\infty) \xrightarrow{\delta} \Omega^1_{\mathbb{P}} \otimes \left(F^{0}_{\mathrm{Del}}\widetilde{\mathcal{M}}\right)((i+2)\infty)\right].$$

Then we have the filtration

$$\widetilde{F_{\mathrm{Del}}}^{-i-1} = \widetilde{F}^{-2i-3} \supset \widetilde{F}^{-2i-2} \supset \widetilde{F}^{-2i-1} = \widetilde{F_{\mathrm{Del}}}^{-i} \quad (i \geq 0)$$

and on each graded piece the map δ is simply the multiplication by dt, which is clearly an isomorphism. □

Theorem 6.2 ([25, Thm.6.1]) *Assume that $(\mathcal{M}, F, \theta)$ is a variation of (pure) Hodge structure on $\mathbb{P}$. With notations as above, we have*

(i) $\mathbb{H}^i\left(\mathbb{P}, \widetilde{\mathcal{M}} \xrightarrow{\delta} \Omega^1 \otimes \widetilde{\mathcal{M}}\right) = 0$ *if* $i \neq 1$, *and*

(ii) *the natural map*

$$\mathbb{H}^i\left(\mathbb{P}, F^{\lambda}_{\mathrm{Del}}\widetilde{\mathcal{M}} \xrightarrow{\delta} \Omega^1 \otimes F^{\lambda-1}_{\mathrm{Del}}\widetilde{\mathcal{M}}\right) \to \mathbb{H}^i\left(\mathbb{P}, \widetilde{\mathcal{M}} \xrightarrow{\delta} \Omega^1 \otimes \widetilde{\mathcal{M}}\right)$$

is injective for any i, λ.

Remark. (i) The above discussion is a special case of the consideration in [25] where the whole construction is carried out in the case that $\mathcal{M}$ is the minimal extension of a variation of Hodge structure on an open dense subset of $\mathbb{P}$. In this case, the definition of $F^{\lambda}_{\mathrm{Del}}\widetilde{\mathcal{M}}$ involves the V-filtration on $\mathcal{M}$ across the singularity. However in our restricted setting, the smoothness of $\mathcal{M}$ on $\mathbb{P}$ implies that $V^{\gamma-\lceil\gamma\rceil}\mathcal{M} = \mathcal{M}$. Thus the definition in [25, §6.b] reduces to (39).

(ii) To be precise, Sabbah considers the twist $\theta - dt$ on $\mathcal{M}$, not $\theta + dt$. Thus one should apply the results in [25] to the variation $\iota^*\mathcal{M}$ where ι is the multiplication by -1 on $\mathbb{P}$. By translating [25, Thm.6.1] applied to $\iota^*\mathcal{M}$ via the isomorphism ι, one obtains the above theorem.

(*b*) *The case of a smooth pencil*

Let $f : X \to \mathbb{P}$ be a smooth proper morphism. Let $U = f^{-1}(\mathbb{A})$ and consider the irregular connection $\nabla = \nabla_f$ on U. In this case, we have $P = S$ and $\mathcal{O}_X(S) = f^*\mathcal{O}_{\mathbb{P}}([\infty])$.

For any integer λ, let

$$\begin{aligned} G^{\lambda} &:= \left[\Omega^{\lambda}_X(S) \xrightarrow{\nabla} \Omega^{\lambda+1}_X(3S) \to \cdots \to \Omega^{\lambda+j}_X((2j+1)S) \to \cdots\right][-\lambda] \\ G^{\lambda}_{\mathrm{rel}} &:= \left[\Omega^{\lambda}_{X/\mathbb{P}}(S) \xrightarrow{d} \Omega^{\lambda+1}_{X/\mathbb{P}}(3S) \to \cdots \to \Omega^{\lambda+j}_{X/\mathbb{P}}((2j+1)S) \to \cdots\right][-\lambda]. \end{aligned}$$

Notice the different chain maps ∇ and d in the definitions. As in the untwisted case, one has the exact sequence of complexes on X

$$0 \to f^*\Omega^1_{\mathbb{P}} \underset{\mathcal{O}_X}{\otimes} G^{\lambda-1}_{\mathrm{rel}}[-1] \to G^{\lambda} \to G^{\lambda}_{\mathrm{rel}} \to 0$$

for every λ since f is smooth and $df = f^*dt$, which is zero in the relative differential forms. Thus on $\mathbb{P}$ one obtains the distinguished triangle

$$\mathcal{R}f_*G^{\lambda}_{\mathrm{rel}} \xrightarrow{\delta} \Omega^1_{\mathbb{P}} \underset{\mathcal{O}_{\mathbb{P}}}{\otimes} \mathcal{R}f_*G^{\lambda-1}_{\mathrm{rel}} \to \mathcal{R}f_*G^{\lambda}[1] \xrightarrow{+1} . \tag{43}$$

We now consider the following relative cohomology on $\mathbb{P}$

$$\begin{aligned} \mathcal{M}_i &:= \mathcal{R}^i f_*\mathbb{C} = \mathcal{R}^i f_* G^0_{\mathrm{rel}} \\ \widetilde{\mathcal{M}}_i &:= \mathcal{M}_i(*\infty) \end{aligned}$$

equipped with the Hodge filtration and the Gauss-Manin connection θ. One then can consider the twisted connection $\delta = \theta + dt$ on $\widetilde{\mathcal{M}}_i$ as in the previous subsection.

Lemma 6.3 *We have*

$$F^\lambda_{\mathrm{Del}}\widetilde{\mathcal{M}}_i = \mathcal{R}^i f_* G^\lambda_{\mathrm{rel}}$$

and under this identification, the connecting map δ *in* (43) *equals the twisted connection defined in* (40) *for the variation* $\mathcal{M} = \mathcal{M}_i$ *of pure Hodge structure.*

Proof. Combining the usual Hodge filtration and the projection formula, we have

$$\left(F^j\mathcal{M}_i\right)(n\infty) = \mathcal{R}^i f_*\left[\Omega^j_{X/\mathbb{P}}(nS) \xrightarrow{d} \Omega^{j+1}_{X/\mathbb{P}}(nS) \to \cdots\right][-j],$$

regarded as a submodule of $\widetilde{\mathcal{M}}$. By the explicit description (41) of the filtration F_{Del}, the first assertion follows.

The second assertion is clear since the differential of the complex G^λ setting in the middle of the commutative diagram (40) is simply given by the twisted connection $\nabla = d + f^*dt$. ☐

Lemma 6.4 *The distinguished triangle* (43) *splits in the sense that for any* i*, one has the quasi-isomorphism*

$$\operatorname{tot}\left[\begin{tikzcd} \mathcal{R}^{i-1}f_*\Big(\Omega^\lambda_{X/\mathbb{P}}(S) \xrightarrow{d} \Omega^{\lambda+1}_{X/\mathbb{P}}(3S) \to \cdots\Big) \arrow[d, "\delta"] \\ \Omega^1_{\mathbb{P}} \otimes \mathcal{R}^i f_*\Big(\underset{\blacktriangle}{\Omega^{\lambda-1}_{X/\mathbb{P}}}(S) \xrightarrow{d} \Omega^\lambda_{X/\mathbb{P}}(3S) \to \cdots\Big) \end{tikzcd}\right] \xrightarrow{\sim}$$

$$\mathcal{R}^i f_*\Big(\Omega^\lambda_X(S) \xrightarrow{\nabla} \Omega^{\lambda+1}_X(3S) \to \cdots\Big).$$

Proof. This is equivalent to show that the upper row in the commutative diagram

$$\begin{tikzcd} 0 \arrow[r] & \mathcal{R}^i f_* G^\lambda_{\mathrm{rel}} \arrow[r, "\delta"] \arrow[d, hook] & \Omega^1_{\mathbb{P}} \otimes \mathcal{R}^i f_* G^{\lambda-1}_{\mathrm{rel}} \arrow[r] \arrow[d, hook] & \mathcal{R}^{i+1} f_* G^\lambda \arrow[r] & 0 \\ & \mathcal{R}^i f_* G^0_{\mathrm{rel}} \arrow[r, "\delta"] & \Omega^1_{\mathbb{P}} \otimes \mathcal{R}^i f_* G^{-1}_{\mathrm{rel}} \arrow[r, equal] & \Omega^1_{\mathbb{P}}(2\infty) \otimes \mathcal{R}^i f_* G^0_{\mathrm{rel}} & \end{tikzcd}$$

is exact for any i, which is enough to show that the connection δ in the upper row is injective. Notice that the injective vertical maps follow from the previous lemma. Now the map δ in the lower row is injective as mentioned in the previous subsection (before Prop.6.1). ☐

The relation with our irregular Hodge filtration $F^\lambda(\nabla)$ on X attached to $\nabla = \nabla_f$ is given in the following.

Lemma 6.5 *For any integer* λ*, the inclusion* $F^\lambda(\nabla) \to G^\lambda$ *of complexes on* X *is a quasi-isomorphism.*

Proof. One only needs to look at the stacks at points in S. Consider the following refinement

$$R(k) := G^{\lambda} \cap \left(\left[\breve{\Omega}^{\lambda}(kS) \xrightarrow{\nabla} \breve{\Omega}^{\lambda+1}((k+1)S) \to \cdots \to \breve{\Omega}^{\lambda+j}((k+j)S) \to \cdots \right] [-\lambda] \right).$$

We have (strict inclusions)

$$F^{\lambda}(\nabla) = R(0) \subset R(1) \subset \cdots \subset R(n-\lambda+1) = G^{\lambda}.$$

We now show that each inclusion above is a quasi-isomorphism.

Let $\{z, x_1, \cdots, x_m\}$ be local coordinates at a point in S such that $f = 1/z$. Then the quotient $Q(k) := R(k+1)/R(k)$ is an $\bar{\mathcal{O}} := \mathcal{O}/z\mathcal{O}$-linear complex with $\nabla =$ left multiplication by $-dz/z^2$. Let us check that the complex $Q(k)$ is exact at degree $\lambda + j$.

When $k = j$, the $\bar{\mathcal{O}}$-module $Q(k)^{\lambda+j}$ is generated by

$$\frac{1}{z^{2j+1}} \bigwedge^{\lambda+j} \{dx_1, \cdots, dx_m\}.$$

It is then obvious that $\nabla : Q(k)^{\lambda+j} \to Q(k)^{\lambda+j+1}$ is injective.

When $k \neq j$, the $\bar{\mathcal{O}}$-module $Q(k)^{\lambda+j}$ is generated by

$$\frac{1}{z^{2j+1}} \bigwedge^{\lambda+j} \{dx_1, \cdots, dx_m\} \quad \text{and} \quad B := \frac{dz}{z^{2j+1}} \bigwedge^{\lambda+j-1} \{dx_1, \cdots, dx_m\}.$$

Thus $\ker \left\{ \nabla : Q(k)^{\lambda+j} \to Q(k)^{\lambda+j+1} \right\}$ is generated by the collection B over $\bar{\mathcal{O}}$, which clearly lies in the image $\nabla \left(Q(k)^{\lambda+j-1} \right)$ of ∇. ☐

Theorem 6.6 *Let $f : X \to \mathbb{P}$ be a proper smooth morphism, $U = f^{-1}(\mathbb{A})$ and $\nabla = \nabla_f$ the irregular connection on U. Then the Hodge to de Rham spectral sequence associated with the irregular Hodge filtration $F^{\lambda}(\nabla)$ degenerates at the initial stage.*

Proof. Notice that in this case the jumps λ in the filtration $F^{\bullet}(\nabla)$ are all integers. We have to show that $H^i\left(U, F^{\lambda}(\nabla)\right) \to H^i\left(U, F^0(\nabla)\right)$ is always an injection for any i and any $\lambda \geq 0$. First by Lemma 6.5, we have $H^i\left(U, F^{\lambda}(\nabla)\right) \xrightarrow{\sim} \mathbb{H}^i\left(X, G^{\lambda}\right)$.

Now consider the commutative diagram

$$\begin{tikzcd}
\mathbb{H}^{j+1}\left(\mathbb{P}, F^{\lambda}_{\mathrm{Del}}\widetilde{\mathcal{M}}_i \xrightarrow{\delta} \Omega^1 \otimes F^{\lambda-1}_{\mathrm{Del}}\widetilde{\mathcal{M}}_i\right) \arrow[r] \arrow[d, "a_{\lambda}"', "\cong"] & \mathbb{H}^{j+1}\left(\mathbb{P}, F^{0}_{\mathrm{Del}}\widetilde{\mathcal{M}}_i \xrightarrow{\delta} \Omega^1 \otimes F^{-1}_{\mathrm{Del}}\widetilde{\mathcal{M}}_i\right) \arrow[d, "\cong"', "a_0"] \\
\mathbb{H}^{j+1}\left(\mathbb{P}, \mathcal{R}^i f_* G^{\lambda}_{\mathrm{rel}} \xrightarrow{\delta} \Omega^1 \otimes \mathcal{R}^i f_* G^{\lambda-1}_{\mathrm{rel}}\right) \arrow[r] \arrow[d, "b_{\lambda}"', "\cong"] & \mathbb{H}^{j+1}\left(\mathbb{P}, \mathcal{R}^i f_* G^{0}_{\mathrm{rel}} \xrightarrow{\delta} \Omega^1 \otimes \mathcal{R}^i f_* G^{-1}_{\mathrm{rel}}\right) \arrow[d, "\cong"', "b_0"] \\
\mathbb{H}^{j}\left(\mathbb{P}, \mathcal{R}^{i+1} f_* G^{\lambda}\right) \arrow[r] & \mathbb{H}^{j}\left(\mathbb{P}, \mathcal{R}^{i+1} f_* G^{0}\right).
\end{tikzcd}$$

Here the horizontal arrows are induced by the natural inclusions of complexes, the two vertical isomorphisms a_{λ}, a_0, are given by Lemma 6.3 and the last two isomorphisms b_{λ}, b_0

are given by Lemma 6.4. By Thm.6.2, the upper arrow is always injective and the two spaces in the upper corners are non-zero only if $j = 0$. Thus the same statements hold for the lower arrow and the lower corner spaces.

It follows that the spectral sequence

$$E_2^{pq} = \mathbb{H}^p\left(\mathbb{P}, \mathcal{R}^q f_* G^\lambda\right) \Longrightarrow \mathbb{H}^{p+q}\left(X, G^\lambda\right)$$

degenerates and yields

$$\mathbb{H}^q\left(X, G^\lambda\right) \xrightarrow{\sim} \mathbb{H}^0\left(\mathbb{P}, \mathcal{R}^q f_* G^\lambda\right).$$

This completes the proof. □

A Comparison with Deligne's definition

In this appendix we recall Deligne's definition of the irregular Hodge filtration in the curve case in [9] and show that it induces the same filtration as ours in the de Rham cohomology.

Consider the pair (U, f) where U is a smooth curve. Let $\nabla = d + df$ be the associated exponentially twisted connection on U as before. Let X be the smooth completion of U with boundary $S := X \setminus U$. Again write $P =$ the pole divisor of f on X.

Deligne then defines inductively an exhaustive and separated filtration $\mathfrak{F}^\lambda$ of the two-term complex $(\Omega_X^\bullet(*S), \nabla)$ by letting

$$\mathfrak{F}^\lambda(\Omega_X^\bullet(*S), \nabla) = \left[\mathfrak{F}^\lambda \mathcal{O}_X(*S) \xrightarrow{\nabla} \mathfrak{F}^\lambda \Omega_X^1(*S)\right]$$

where

$$\begin{aligned}
\mathfrak{F}^\lambda \mathcal{O}_X(*S) &= \begin{cases} 0 & \text{if } \lambda > 0 \\ \mathcal{O}_X(S - \lceil \lambda P \rceil) & \text{if } -1 < \lambda \le 0 \\ \left(\mathfrak{F}^{\lambda+1}\mathcal{O}_X(*S)\right)(S+P) & \text{if } \lambda \le -1 \end{cases} \\
\mathfrak{F}^\lambda \Omega_X^1(*S) &= \Omega_X^1 \underset{\mathcal{O}_X}{\otimes} \left(\mathfrak{F}^{\lambda-1}\mathcal{O}_X(*S)\right).
\end{aligned}$$

Define a subcomplex $\Omega_X^\bullet(\log_{\mathfrak{F}} \nabla)$ of $(\Omega_X^\bullet(*S), \nabla)$ to be the two-term complex

$$\left[\ \ker\left\{\mathfrak{F}^0 \mathcal{O}_X(*S) \xrightarrow{\nabla} \mathrm{Gr}_{\mathfrak{F}}^0 \Omega_X^1(*S)\right\} \xrightarrow{\nabla} \mathfrak{F}^{0+}\Omega_X^1(*S)\ \right]$$

equipped with the induced filtration $\mathfrak{F}^\lambda$. Then $\mathfrak{F}^\lambda \Omega_X^\bullet(\log_{\mathfrak{F}} \nabla)$ is non-trivial only if $0 \le \lambda \le 1$. We call $\Omega_X^\bullet(\log_{\mathfrak{F}} \nabla)$ the logarithmic subcomplex of $(\Omega_X^\bullet(*S), \nabla)$; it is a complex of coherent sheaves on X, filtered by coherent subcomplexes. The context of the irregular Hodge theory over curves is then summarized as the following.

Theorem A.1 (Deligne) *With notations as above, we have the following.*

(i) *The natural inclusion* $(\Omega_X^\bullet(\log_{\mathfrak{F}} \nabla), \nabla, \mathfrak{F}) \to (\Omega_X^\bullet(*S), \nabla, \mathfrak{F})$ *is a quasi-isomorphism of filtered complexes on* X.

(ii) *For each* λ*, the map* $\mathbb{H}(X, \mathfrak{F}^\lambda) \to H_{\mathrm{dR}}(U, \nabla)$ *induced by the inclusion of complexes is injective (i.e. the spectral sequence associated with the filtration* $\mathfrak{F}$ *degenerates at the initial* E_1 *stage).*

We remark again that the construction can be generalized to exponential twists of unitary regular connections of any ranks over the curve U and the corresponding statements as above continue to hold in the general case.

Now let us compare the two filtrations $F^\bullet(\nabla)$ and $\mathfrak{F}^\bullet$. First we clearly have $F^\lambda(\nabla) \subset \mathfrak{F}^\lambda$ for any $\lambda \in \mathbb{R}$. On the other hand, one readily observes that the two corresponding logarithmic filtered complexes $\Omega^\bullet_X(\log_F \nabla)$ and $\Omega^\bullet_X(\log_{\mathfrak{F}} \nabla)$ are exactly the same subcomplex of $(\Omega^\bullet_X(*S), \nabla)$. Thus we obtain the following statement.

Proposition A.2 *In the curve case, the two filtrations* $F^\bullet(\nabla)$ *and* $\mathfrak{F}^\bullet$ *induce the same filtration on the twisted de Rham cohomology* $H_{\mathrm{dR}}(U, \nabla)$*.*

References

[1] A. Adolphson and S. Sperber, Exponential sums and Newton polyhedra: cohomology and estimates. *Ann. of Math. (2)* 130 (1989), no. 2, 367-406.

[2] A. Adolphson and S. Sperber, On twisted de Rham cohomology. *Nagoya Math. J.* 146 (1997), 55-81.

[3] F. Baldassarri and P. Berthelot, On Dwork cohomology for singular hypersurfaces. *Geometric aspects of Dwork theory. Vol. I, II*, 177-244, Walter de Gruyter GmbH & Co. KG, Berlin, 2004.

[4] W. Barth, C. Peters and A. Van de Ven, *Compact complex surfaces.* Ergebnisse der Mathematik und ihrer Grenzgebiete (3), 4. Springer-Verlag, Berlin, 1984.

[5] P. Berthelot and A. Ogus, *Notes on crystalline cohomology.* Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1978.

[6] S. Bloch and H. Esnault, Homology for irregular connections. *J. Théor. Nombres Bordeaux* 16 (2004), no. 2, 357-371.

[7] P. Deligne, *Équations différentielles à points singuliers réguliers.* Lecture Notes in Mathematics, 163. Springer-Verlag, Berlin-New York, 1970.

[8] P. Deligne and L. Illusie, Relèvements modulo p^2 et décomposition du complexe de de Rham. *Invent. Math.* 89 (1987), no. 2, 247-270.

[9] P. Deligne, B. Malgrange and J.-P. Ramis, *Singularités irrégulières. Correspondance et documents.* Documents Mathématiques, 5. Société Mathématique de France, Paris, 2007.

[10] F. El Zein, *Introduction à la théorie de Hodge mixte.* Actualités Mathématiques. Hermann, Paris, 1991.

[11] W. Fulton, *Introduction to toric varieties.* Annals of Mathematics Studies, 131. Princeton University Press, Princeton, NJ, 1993.

[12] R. Godement, *Topologie algébrique et théorie des faisceaux.* Troisième édition revue et corrigée. Publications de l'Institut de Mathématique de l'Université de Strasbourg, XIII. Actualités Scientifiques et Industrielles, No. 1252. Hermann, Paris, 1973.

[13] R. Hartshorne, *Residues and duality.* Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics, 20. Springer-Verlag, Berlin-New York 1966.

[14] R. Hartshorne, *Algebraic geometry.* Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977.

[15] M. Hien, Periods for flat algebraic connections. *Invent. Math.* 178 (2009), no. 1, 1-22.

[16] R. Hotta, K. Takeuchi and T. Tanisaki, *D-modules, perverse sheaves, and representation theory.* Progress in Mathematics, 236. Birkhäuser Boston, Inc., Boston, MA, 2008.

[17] L. Illusie, Complexe de de Rham-Witt et cohomologie cristalline. *Ann. Sci. École Norm. Sup. (4)* 12 (1979), no. 4, 501-661.

[18] L. Katzarkov, M. Kontsevich and T. Pantev, Hodge theoretic aspects of mirror symmetry. *From Hodge theory to integrability and TQFT tt*-geometry*, 87-174, Proc. Sympos. Pure Math., 78, Amer. Math. Soc., Providence, RI, 2008.

[19] M. Kontsevich and Y. Soibelman, Cohomological Hall algebra, exponential Hodge structures and motivic Donaldson-Thomas invariants. *Commun. Number Theory Phys.* 5 (2011), no. 2, 231-352.

[20] A. G. Kouchnirenko, Polyèdres de Newton et nombres de Milnor. *Invent. Math.* 32 (1976), no. 1, 1-31.

[21] N. O. Nygaard, Slopes of powers of Frobenius on crystalline cohomology. *Ann. Sci. École Norm. Sup. (4)* 14 (1981), no. 4, 369-401 (1982).

[22] A. Ogus, F-crystals, Griffiths transversality, and the Hodge decomposition. *Astérisque* No. 221 (1994).

[23] A. Ogus and V. Vologodsky, Nonabelian Hodge theory in characteristic p. *Publ. Math. Inst. Hautes Études Sci.* No. 106 (2007), 1-138.

[24] C. A. M. Peters and J. H. M. Steenbrink, *Mixed Hodge structures.* Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, 52. Springer-Verlag, Berlin, 2008.

[25] C. Sabbah, Fourier-Laplace transform of a variation of polarized complex Hodge structure, II. *New developments in algebraic geometry, integrable systems and mirror symmetry (RIMS, Kyoto, 2008)*, 289-347, Adv. Stud. Pure Math., 59, Math. Soc. Japan, Tokyo, 2010.

[26] C. Sabbah, Théorie de Hodge et correspondence de Hitchin-Kobayashi sauvages [d'après T. Mochizuki]. arXiv:1201.4531.

[27] D. Wan, Variation of p-adic Newton polygons for L-functions of exponential sums. *Asian J. Math.* 8 (2004), no. 3, 427-471.

[28] J. Włodarczyk, Simple constructive weak factorization. *Algebraic geometry – Seattle 2005. Part 2*, 957-1004, Proc. Sympos. Pure Math., 80, Part 2, Amer. Math. Soc., Providence, RI, 2009.

[29] H. J. Zhu, L-functions of exponential sums over one-dimensional affinoids: Newton over Hodge. *Int. Math. Res. Not.* 2004, no. 30, 1529-1550.